

10/616,390

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
20 December 2001 (20.12.2001)

PCT

(10) International Publication Number  
**WO 01/96584 A2**(51) International Patent Classification<sup>7</sup>: **C12N 15/82**(21) International Application Number: **PCT/US01/18911**(22) International Filing Date: **12 June 2001 (12.06.2001)**(25) Filing Language: **English**(26) Publication Language: **English**(30) Priority Data:  
60/210,917 **12 June 2000 (12.06.2000) US**(71) Applicant (for all designated States except US):  
**AKKADIX CORPORATION [US/US]; 4204 Sorrento Valley Blvd., Suite A, San Diego, CA 92121-1412 (US).**


(72) Inventors; and

(75) Inventors/Applicants (for US only): **MUSHEGIAN, Arcady, R. [—/US]; 3987 Santa Nella Place, San Diego, CA 92130 (US). TAYLOR, Christopher, G. [—/US]; 2910-A**Luciema Street, Carlsbad, CA 92009 (US). **FEITELSON, Jerald, S. [—/US]; 4387 Mistral Place, San Diego, CA 92130 (US). EROSHKIN, Alexy, M. [—/US]; 3803 Ruelle San Rafael, San Diego, CA 92130 (US).**(74) Agents: **LLOYD, Jeff et al.; Saliwanchik, Lloyd & Saliwanchik, Suite A-1, 2421 N.W. 41st Street, Gainesville, FL 32606 (US).**

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: **MATERIALS AND METHODS FOR THE CONTROL OF NEMATODES**

(57) Abstract: The subject invention provides novel methods and compositions for controlling nematodes. More specifically, the subject invention provides RNAi molecules, polynucleotide sequences, and methods of using these sequences in nematode control.

WO 01/96584 A2



**Published:**

— without international search report and to be republished  
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## DESCRIPTION

### MATERIALS AND METHODS FOR THE CONTROL OF NEMATODES

#### Background of the Invention

[0001] Plant parasitic nematodes, such as root-knot nematodes (*Meloidogyne* species) and cyst nematodes (*Globodera* and *Heterodera*), attack nearly every food crop, and are among the world's most damaging agricultural pests. For example, root-knot nematodes parasitize more than 2,000 plant species from diverse plant families and represent a tremendous threat to crop production world-wide. These biotrophic pathogens have evolved highly specialized and complex feeding relationships with their hosts.

[0002] Nematodes cause millions of dollars of damage each year to turf grasses, ornamental plants, and food crops. Efforts to eliminate or minimize damage caused by nematodes in agricultural settings have typically involved the use of soil fumigation with materials such as chloropicrin, methyl bromide, and dazomet, which volatilize to spread the active ingredient throughout the soil. Such fumigation materials can be highly toxic and may create an environmental hazard. Various non-fumigant chemicals have also been used, but these too create serious environmental problems and can be highly toxic to humans.

[0003] Some research articles have been published concerning the effects of  $\delta$ -endotoxins from *B. thuringiensis* species on the viability of nematodes. See, for example, Bottjer, Bone and Gill ([1985] *Experimental Parasitology* 60:239-244); Ignoffo and Dropkin (Ignoffo, C.M., Dropkin, V.H. [1977] *J. Kans. Entomol. Soc.* 50:394-398); and Ciordia, H. and W.E. Bizzell ([1961] *Jour. of Parasitology* 47:41 [abstract]). Several patents have issued describing the control of nematodes with *B.t.* See, for example, U.S. Patent Nos. 4,948,734; 5,093,120; 5,281,530; 5,426,049; 5,439,881; 5,236,843; 5,322,932; 5,151,363; 5,270,448; 5,350,577; 5,667,993; and 5,670,365. The development of resistance by insects to *B.t.* toxins is one obstacle to the successful use of such toxins.

[0004] The pesticidal activity of avermectins is well known. The avermectins are disaccharide derivatives of pentacyclic, 16-membered lactones. They can be divided into four major compounds: A<sub>1a</sub>, A<sub>2a</sub>, B<sub>1a</sub>, and B<sub>2a</sub>; and four minor compounds: A<sub>1b</sub>, A<sub>2b</sub>, B<sub>1b</sub>, and B<sub>2b</sub>. The isolation and purification of these compounds is also described in U.S. Patent No. 4,310,519, issued January 12, 1982. Avermectin B<sub>2a</sub> is active against the root-knot nematode, *Meloidogyne incognita*. It is reported to be 10-30 times as potent as commercial contact nematicides when incorporated into soil at 0.16-0.25 kg/ha (Boyce Thompson Institute for Plant Research 58th Annual Report [1981]; Putter, L. *et al.* [1981] "Avermectins: Novel Insecticides, Acaracides, and Nematicides from a Soil Microorganism," *Experientia* 37:963-964). Avermectin B<sub>2a</sub> is not toxic to tomatoes or cucumbers at rates of up to 10 kg/ha.

[0005] Fatty acids are a class of natural compounds which occur abundantly in nature and which have interesting and valuable biological activities. Tarjan and Cheo (Tarjan, A.C., P.C. Cheo [1956] "Nematocidal Value of Some Fatty Acids," Bulletin 332, Contribution 884, Agricultural Experiment Station, University of Rhode Island, Kingston, 41 pp.) report the activity of certain fatty acids against nematodes. In 1977 Sitaramaiah and Singh (Sitaramaiah, K., R.S. Singh [1977] *Indian J. Nematol.* 7:58-65) also examined the response of nematodes to fatty acids. The results of these tests with short chain acids were equivocal, showing nematode-inhibitory action in some instances and stimulatory activity in other instances. Phytotoxicity of these acids was observed at higher concentrations. The short chain fatty acids were also examined by Malik and Jairajpuri (Malik, Z., M.S. Jairajpuri [1977] *Nematol. medit.* 12:73-79), who observed nematode toxicity at high concentrations of the fatty acids.

[0006] Notwithstanding the foregoing (some of the limitations of and problems associated with these approaches are discussed above), there is a need for safe and effective alternatives for controlling nematodes.

[0007] One method for disrupting normal cellular processes is by the use double-stranded interfering RNA (RNAi), or RNA-mediated interference (RNAi). When RNAi corresponding to a sense and antisense sequence of a target mRNA is introduced into a cell, the targeted mRNA is degraded and protein translation of that message is stopped. Although not yet fully understood, the mechanism of this post-transcriptional gene



silencing appears to be at least partially due to the generation of small RNA molecules, about 21 - 25 nucleotides in length, that correspond to the sense and antisense pieces of the RNAi introduced into the cell (Bass, B. L. [2000] "Double-stranded RNA as a template for gene silencing" *Cell* 101:235-238).

[0008] The specificity of this gene silencing mechanism appears to be extremely high, blocking expression only of targeted genes, while leaving other genes unaffected. A recent example of the use of RNAi; to inhibit genetic function in plants used *Agrobacterium tumefaciens*-mediated transformation of *Arabidopsis thaliana* (Chuang, C.-F. and E. M. Meyerowitz [2000] "Specific and heritable genetic interference by double-stranded RNA in *Arabidopsis thaliana*" *Proc. Natl. Acad. Sci. USA* 97:4985-4990). Chuang *et al.* describe the construction of vectors delivering variable levels of RNAi targeted to each of four genes involved in floral development. Severity of abnormal flower development varied between transgenic lines. For one of the genes, AGAMOUS (AG), a strong correlation existed between declining accumulation of mRNA and increasingly severe phenotypes, suggesting that AG-specific endogenous mRNA is the target of RNAi.

#### Brief Summary of the Invention

[0009] The subject invention provides novel methods and compositions for controlling nematodes. More specifically, the subject invention provides polynucleotide sequences that encode nematode genes, RNAi that selectively targets mRNA transcripts of these essential nematode genes, and methods of using these sequences in nematode control strategies. Such sequences for use according to the subject invention are summarized in Appendix 1. RNAi molecules disclosed herein can be used to inhibit the expression of one or more of these genes in nematodes.

### Brief Description of the Drawings

[00010] **Figure 1:** Modular Binary Construct System (MBCS): A series of six, 8-base cutter restriction enzyme sites has been placed between the left and right Ti borders of a previously created kan<sup>R</sup>/tet<sup>R</sup> binary plasmid.

[00011] **Figure 2:** An exemplary shuttle vector created for cloning of useful DNA fragments by containing the multi-cloning site (MCS) of a modified Bluescript plasmid flanked by 8-base restriction sites.

[00012] **Figure 3:** An exemplary shuttle vector with exemplary inserts.

[00013] **Figure 4:** A suggested RNAi binary vector with exemplary inserts.

[00014] **Figure 5:** Exemplary selectable markers for MBCS.

[00015] **Figure 6:** Exemplary scorable markers for MCBS.

[00016] **Figure 7:** Exemplary RNAi binary vector.

[00017] **Figure 8:** Exemplary RNAi shuttle vector.

### Brief Description of the Sequences

[00018] Brief Description of the Sequences can be found in Appendix I.

### Detailed Disclosure of the Invention

[00019] The subject invention provides novel methods and compositions for controlling nematodes. More specifically, the subject invention provides polynucleotide sequences and methods of using these sequences in nematode control strategies. A preferred method for controlling nematodes according to the subject invention provides materials and methods for controlling nematodes by using double-stranded interfering RNA (RNAi), or RNA-mediated interference (RNAi). The terms RNAi and RNAi are used interchangeably herein unless otherwise noted.

[00020] In one embodiment of the invention, RNAi molecules are provided which are useful in methods of killing nematodes and/or inhibiting their growth, development, parasitism or reproduction. RNAi molecules of the invention are also useful for the regulation of levels of specific mRNA in nematodes.

[00021] dsRNA (RNAi) typically comprises a polynucleotide sequence identical to a target gene (or fragment thereof) linked directly, or indirectly, to a polynucleotide

sequence complementary to the sequence of the target gene (or fragment thereof). The dsRNA may comprise a polynucleotide linker (stuffer) sequence of sufficient length to allow for the two polynucleotide sequences to fold over and hybridize to each other; however, a linker sequence is not necessary. The linker (stuffer) sequence is designed to separate the antisense and sense strands of RNAi significantly enough to limit the effects of steric hindrances and allow for the formation of dsRNA molecules.

[00022] RNA containing a nucleotide sequence identical to a fragment of the target gene is preferred for inhibition; however, RNA sequences with insertions, deletions, and point mutations relative to the target sequence can also be used for inhibition. Sequence identity may be optimized by sequence comparison and alignment algorithms known in the art (see Gribskov and Devereux, *Sequence Analysis Primer*, Stockton Press, 1991, and references cited therein) and calculating the percent difference between the nucleotide sequences by, for example, the Smith-Waterman algorithm as implemented in the BESTFIT software program using default parameters (e.g., University of Wisconsin Genetic Computing Group). Alternatively, the duplex region of the RNA may be defined functionally as a nucleotide sequence that is capable of hybridizing with a fragment of the target gene transcript.

[00023] As disclosed herein, 100% sequence identity between the RNA and the target gene is not required to practice the present invention. Thus the invention has the advantage of being able to tolerate sequence variations that might be expected due to genetic mutation, strain polymorphism, or evolutionary divergence.

[00024] RNA may be synthesized either *in vivo* or *in vitro*. Endogenous RNA polymerase of the cell may mediate transcription *in vivo*, or cloned RNA polymerase can be used for transcription *in vivo* or *in vitro*. For transcription from a transgene *in vivo* or an expression construct, a regulatory region (e.g., promoter, enhancer, silencer, splice donor and acceptor, polyadenylation) may be used to transcribe the RNA strand (or strands). Inhibition may be targeted by specific transcription in an organ, tissue, or cell type; stimulation of an environmental condition (e.g., infection, stress, temperature, chemical inducers); and/or engineering transcription at a developmental stage or age. The RNA strands may or may not be polyadenylated; the RNA strands may or may not be capable of being translated into a polypeptide by a cell's translational apparatus. RNA

may be chemically or enzymatically synthesized by manual or automated reactions. The RNA may be synthesized by a cellular RNA polymerase or a bacteriophage RNA polymerase (e.g., T3, T7, SP6). The use and production of an expression construct are known in the art (see, for example, WO 97/32016; U.S. Pat. Nos. 5,593,874; 5,698,425; 5,712,135; 5,789,214; and 5,804,693; and the references cited therein). If synthesized chemically or by *in vitro* enzymatic synthesis, the RNA may be purified prior to introduction into the cell. For example, RNA can be purified from a mixture by extraction with a solvent or resin, precipitation, electrophoresis, chromatography, or a combination thereof. Alternatively, the RNA may be used with no or a minimum of purification to avoid losses due to sample processing. The RNA may be dried for storage or dissolved in an aqueous solution. The solution may contain buffers or salts to promote annealing, and/or stabilization of the duplex strands.

[00025] Preferably and most conveniently, RNAi can be targeted to an entire polynucleotide sequence of a gene set forth herein. Preferred RNAi molecules of the instant invention are highly homologous or identical to the polynucleotides summarized in Appendix 1. The homology is preferably greater than 90% and is most preferably greater than 95%.

[00026] Fragments of genes can also be targeted. These fragments are typically in the approximate size range of about 20 nucleotides. Thus, targeted fragments are preferably at least about 15 nucleotides. In certain embodiments, the gene fragment targeted by the RNAi molecule is about 20-25 nucleotides in length. However, other size ranges can also be used. For example, using a *C. elegans* microinjection assay, RNAi "fragments" of about 60 nucleotides with between 95 and 100% identity (to a nematode gene) were determined to cause excellent inhibition.

[00027] Thus, RNAi molecules of the subject invention are not limited to those that are targeted to the full-length polynucleotide or gene. The nematode gene product can be inhibited with a RNAi molecule that is targeted to a portion or fragment of the exemplified polynucleotides; high homology (90-95%) or identity is also preferred, but not necessarily essential, for such applications.

[00028] The polynucleotide sequences identified in Appendix A and shown in the Sequence ID listing are from genes encoding nematode proteins having the functions

shown in Appendix 1. The genes exemplified herein are representative of particular classes of proteins which are preferred targets for disruption according to the subject invention. These classes of proteins include, for example, proteins involved in ribosome assembly; neurot transmitter receptors and ligands; electron transport proteins; metabolic pathway proteins; and protein and polynucleotide production, folding, and processing proteins.

[00029] Genetic regulatory sequences, such as promoters, enhancers, and terminators, can be used in genetic constructs to practice the subject invention. Such constructs themselves can also be used for nematode control. Various constructs can be used to achieve expression in specific plant tissues (by using root specific promoters, for example) and/or to target specific nematode tissues (by using targeting elements or adjacent targeting sequences, for example).

[00030] In a specific embodiment of the subject invention, plant cells, preferably root cells, are genetically modified to produce at least one RNAi that is designed to be taken up by nematodes during feeding to block expression (or the function of) of a target gene. As is known in the art, RNAi can target and reduce (and, in some cases, prevent) the translation of a specific gene product. RNAi can be used to reduce or prevent message translation in any tissue of the nematode because of its ability to cross tissue and cellular boundaries. Thus, RNAi that is contacted with a nematode by soaking, injection, or consumption of a food source will cross tissue and cellular boundaries. RNAi can also be used as an epigenetic factor to prevent the proliferation of subsequent generations of nematodes.

[00031] Nematode polynucleotide sequences disclosed herein demonstrate conserved nucleotide motifs among different nematode genera. Conserved nucleotide motifs strongly suggest that these sequences are associated with viability and/or parasitism and are functionally conserved and expressed in both *Meloidogyne incognita* (root-knot nematode) and *Globodera rostochiensis* and *Globodera pallida* (potato cyst nematodes). The use of these polynucleotides, and RNAi inhibitors thereof, is advantageous because such RNAi can be designed to have broad RNAi specificity and are thus useful for controlling a large number of plant parasitic nematodes *in planta*. Because the genes identified in this disclosure are associated with nematode survival

and/or parasitism, RNAi inhibition of these genes (arising from contacting nematodes with compositions comprising RNAi molecules) prevents and/or reduces parasitic nematode growth, development, and or parasitism.

[00032] Methods of the subject invention include the transformation of plant cells with genes or polynucleotides of the present invention, which can be used to produce nematode inhibitors or RNAi in the plants. In one embodiment, the transformed plant or plant tissue can express RNAi molecules encoded by the gene or polynucleotide sequence introduced into the plant. Other nematode inhibitors contemplated by the invention include antisense molecules specific to the polynucleotide sequences disclosed herein. The transformation of plants with genetic constructs disclosed herein can be accomplished using techniques well known to those skilled in the art and can involve modification of the gene(s) to optimize expression in the plant to be made resistant to nematode infection and infestation. Furthermore, it is known in the art that many tissues of the transgenic plants (such as the roots) can be targeted for transformation.

[00033] RNA-mediated interference (RNAi) of gene expression. Several aspects of root-knot nematode biology make classical genetic studies difficult with this organism. Since root-knot nematodes reproduce by obligatory mitotic parthenogenesis, the opportunity to perform genetic crosses is not available. Microinjection of RNAi can be used to manipulate gene expression in *C. elegans* (Fire, A., S. Xu, M. K. Montgomery, S. A. Kostas, S. E. Driver, and C. C. Mello. [1998] "Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*" *Nature* 391:806- 811). Microinjecting (into adult nematodes) RNAi can turn off specific genes in progeny worms complementary to the coding region of the genes. Moreover, gene inhibition occurs in progeny when RNAi is injected into the body cavity of the adult, indicating the ability of the RNAi to cross cellular boundaries. This RNAi injection method provides a molecular genetic tool that allows for analysis of gene function in root-knot nematodes.

[00034] RNAi can be taken up by *C. elegans* by simply soaking the nematodes in a solution RNAi. This results in targeted inhibition of gene expression in the nematode (Maeda, I., Y. Kohara, M. Yamamoto and A. Sugimoto [1999] "RNAi screening with a non-redundant cDNA set" International Worm Meeting, Madison, WI, abstract 565). Nematodes fed *E. coli* expressing RNAi also demonstrate targeted and

heritable inhibition of gene expression (Sarkissian, M., H. Tabara and C. C. Mello [1999] "A mut-6 screen for RNAi deficient mutants" International Worm Meeting, Madison, WI, abstract 741; Timmons, I. and A. Fire [1998] "Specific interference by ingested dsRNA" *Nature* 395:854; WO 99/32619, hereby incorporated by reference in its entirety).

[00035] Accordingly, one aspect of the instant invention is directed to the control of nematodes comprising contacting nematodes with compositions comprising RNAi molecules specific to the nematode genes disclosed herein. The contacting step may include soaking the nematodes in a solution containing RNAi molecules, feeding nematodes RNAi molecules contained in microbes or plant cells upon which the nematode feeds, or injecting nematodes with RNAi. Nematodes can also be "contacted" and controlled by RNAi expressed in plant tissues that would be consumed, ingested, or frequented by nematodes.

[00036] The RNAi molecules provided to the nematodes may be specific to a single gene. A "cocktail" of RNAi molecules specific to various segments of a single gene can also be used. In addition, a "multigene cocktail" of RNAi molecules specific to two or more genes (or segments thereof) may be applied to the nematodes according to the subject invention.

[00037] In addition to RNAi uptake mediated by transgenic plants, nematodes can be directly transformed with RNAi constructs of cDNAs encoding secretory or other essential proteins to reduce expression of the corresponding gene. The transgenic animals can be assayed for inhibition of gene product using immunoassays or for reduced virulence on a host. Progeny of affected worms can also be assayed by similar methods.

[00038] Procedures that can be used for the preparation and injection of RNAi include those detailed by Fire *et al.*, (1998; <ftp://ciw1.ciwemb.edu>). Root-knot nematodes can be routinely monoxenically cultured on *Arabidopsis thaliana* roots growing on Gamborg's B-5/Gelrite® media. This nematode-host pathosystem is ideally suited for these microinjection experiments since limited root galling results in the parasitic stages (late J2 through adult females) developing outside of the root for easy accessibility for injecting. Another advantage is the parthenogenic reproduction of root-knot nematodes, which makes fertilization by males unnecessary for egg production. The RNAi can be injected into the body cavity of parasitic stages of root-knot nematodes

feeding on *A. thaliana* roots using microinjection. Control nematodes can be injected in parallel with only buffer or an unrelated RNAi. Injected nematodes can be monitored for egg production, and the eggs can be collected for the assays described below. Female root-knot nematodes will typically survive and lay more than 250 eggs following 1  $\mu$ l injection of buffer.

[00039] Alternatively, methods are available for microinjecting materials directly into the plant root cells upon which nematodes feed: giant cells or syncytial cells (Böckenhoff, A. and F.M.W. Grundler [1994] "Studies on the nutrient uptake by the beet cyst nematode *Heterodera schachtii* by *in situ* microinjection of fluorescent probes into the feeding structures in *Arabidopsis thaliana*" *Parasitology* 109:249-254). This provides an excellent test system to screen RNAi molecules for efficacy by directly inhibiting growth and development of the nematode feeding upon the microinjected plant cell, or by reducing fecundity and the ability of said nematode to generate pathogenic or viable progeny.

[00040] There are a number of strategies that can be followed to assay for RNAi gene interference. Inhibition of gene expression by RNAi inhibits the accumulation of the corresponding secretory protein in the esophageal gland cells of transgenic J2 hatched from the eggs produced by the injected nematodes. In the first assay, polyclonal antibodies to the target gene product can be used in immunolocalization studies (Hussey, R. S. [1989] "Monoclonal antibodies to secretory granules in esophageal glands of *Meloidogyne* species" *J. Nematol.* 21:392-398; Borgonie, G, E. van Driessche, C. D. Link, D. de Waele, and A. Coomans [1994] "Tissue treatment for whole mount internal lectin staining in the nematodes *Caenorhabditis elegans*, *Panagrolaimus superbus* and *Acrobeloides maximus*" *Histochemistry* 101:379-384) to monitor the synthesis of the target protein in the gland cells of progeny of the injected nematodes, or in any other nematode tissue that fails to express the essential targeted gene. Interference of endogenous gene activity by the RNAi eliminates binding of the antibodies to secretory granules in the glands, or any other target tissue, of the transgenic nematodes, and can be monitored by these *in situ* hybridization experiments. Control nematodes injected only with the injection buffer can be processed similar to the RNAi treated nematodes.



[00041] Another assay is designed to determine the effect of the RNAi on reducing the virulence of J2 progeny of the injected females. Egg masses from injected females can be transferred singly to *A. thaliana* plates to assess the ability of the transgenic J2 to infect roots. The J2 hatching from the eggs transferred to the plates can be monitored; after 25 days the number of galls with egg laying females can be recorded. The *A. thaliana* roots can also be stained with acid fuchsin to enumerate the number of nematodes in the roots. Egg masses from nematodes injected only with the injection buffer can be handled similarly and used as controls. The treatments can be replicated, and the root infection data can be analyzed statistically. These experiments can be used to assess the importance of the target genes in root-knot nematode's virulence or viability. By staining the J2 progeny of the injected females with the antibodies, it can be determined whether RNAi blocks expression of the targeted gene.

[00042] Additional uses of polynucleotides. The polynucleotide sequences exemplified herein can be used in a variety of ways. These polynucleotides can be used in assays for additional polynucleotides and additional homologous genes, and can be used in tracking the quantitative and temporal expression of parasitism genes in nematodes. These polynucleotides can be cloned into microbes for production and isolation of their gene products. Among the many uses of the isolated gene product is the development of additional inhibitors and modifiers. The protein products of the subject polynucleotides can also be used as diagnostic tools. For example, proteins encoded by the parasitism genes, as identified herein, can be used in large scale screenings for additional peptide inhibitors. The use of peptide phage display screening is one method that can be used in this regard. Thus, the subject invention also provides new biotechnological strategies for managing nematodes under sustainable agricultural conditions.

[00043] Antisense technologies can also be used for phytopathogenic nematode control. Antisense technology can be used to interfere with expression of the disclosed endogenous nematode genes. Antisense technology can also be used to alter the components of plants used as targets by the nematodes. For example, the transformation of a plant with the reverse complement of an endogenous gene encoded by a polynucleotide exemplified herein can result in strand co-suppression and gene silencing

or inhibition of a target involved in the nematode infection process. Thus, the subject invention includes transgenic plants (which are preferably made nematode-resistant in this manner, and other organisms including microbes and phages) comprising RNAi or antisense molecules specific to any of the polynucleotides identified herein.

[00044] Polynucleotide probes. DNA possesses a fundamental property called base complementarity. In nature, DNA ordinarily exists in the form of pairs of anti-parallel strands, the bases on each strand projecting from that strand toward the opposite strand. The base adenine (A) on one strand will always be opposed to the base thymine (T) on the other strand, and the base guanine (G) will be opposed to the base cytosine (C). The bases are held in apposition by their ability to hydrogen bond in this specific way. Though each individual bond is relatively weak, the net effect of many adjacent hydrogen bonded bases, together with base stacking effects, is a stable joining of the two complementary strands. These bonds can be broken by treatments such as high pH or high temperature, and these conditions result in the dissociation, or "denaturation," of the two strands. If the DNA is then placed in conditions which make hydrogen bonding of the bases thermodynamically favorable, the DNA strands will anneal, or "hybridize," and reform the original double-stranded DNA. If carried out under appropriate conditions, this hybridization can be highly specific. That is, only strands with a high degree of base complementarity will be able to form stable double-stranded structures. The relationship of the specificity of hybridization to reaction conditions is well known. Thus, hybridization may be used to test whether two pieces of DNA are complementary in their base sequences. It is this hybridization mechanism which facilitates the use of probes of the subject invention to readily detect and characterize DNA sequences of interest.

[00045] The specifically exemplified polynucleotides of the subject invention can themselves be used as probes. Additional polynucleotide sequences can be added to the ends of (or internally in) the exemplified polynucleotide sequences so that polynucleotides that are longer than the exemplified polynucleotides can also be used as probes. Thus, isolated polynucleotides comprising one or more of the exemplified sequences are within the scope of the subject invention. Polynucleotides that have less nucleotides than the exemplified polynucleotides can also be used and are contemplated within the scope of the present invention. For example, for some purposes, it might be

useful to use a conserved sequence from an exemplified polynucleotide wherein the conserved sequence comprises a portion of an exemplified sequence. Thus, polynucleotides of the subject invention can be used to find additional, homologous (wholly or partially) genes.

[00046] Probes of the subject invention may be composed of DNA, RNA, or PNA (peptide nucleic acid). The probe will normally have at least about 10 bases, more usually at least about 17 bases, and may have about 100 bases or more. Longer probes can readily be utilized, and such probes can be, for example, several kilobases in length. The probe sequence is designed to be at least substantially complementary to a portion of a gene encoding a protein of interest. The probe need not have perfect complementarity to the sequence to which it hybridizes. The probes may be labeled utilizing techniques that are well known to those skilled in this art.

[00047] One approach for the use of the subject invention as probes entails first identifying DNA segments that are homologous with the disclosed nucleotide sequences using, for example, Southern blot analysis of a gene bank. Thus, it is possible, without the aid of biological analysis, to know in advance the probable activity of many new polynucleotides, and of the individual gene products expressed by a given polynucleotide. Such an analysis provides a rapid method for identifying commercially valuable compositions.

[00048] One hybridization procedure useful according to the subject invention typically includes the initial steps of isolating the DNA sample of interest and purifying it chemically. Either lysed nematodes or total fractionated nucleic acid isolated from nematodes can be used. Cells can be treated using known techniques to liberate their DNA (and/or RNA). The DNA sample can be cut into pieces with an appropriate restriction enzyme. The pieces can be separated by size through electrophoresis in a gel, usually agarose or acrylamide. The pieces of interest can be transferred to an immobilizing membrane.

[00049] The particular hybridization technique is not essential to the subject invention. As improvements are made in hybridization techniques, they can be readily applied.

[00050] The probe and sample can then be combined in a hybridization buffer solution and held at an appropriate temperature until annealing occurs. Thereafter, the membrane is washed free of extraneous materials, leaving the sample and bound probe molecules typically detected and quantified by autoradiography and/or liquid scintillation counting. As is well known in the art, if the probe molecule and nucleic acid sample hybridize by forming a strong non-covalent bond between the two molecules, it can be reasonably assumed that the probe and sample are essentially identical or very similar. The probe's detectable label provides a means for determining in a known manner whether hybridization has occurred.

[00051] In the use of the nucleotide segments as probes, the particular probe is labeled with any suitable label known to those skilled in the art, including radioactive and non-radioactive labels. Typical radioactive labels include  $^{32}\text{P}$ ,  $^{35}\text{S}$ , or the like. Non-radioactive labels include, for example, ligands such as biotin or thyroxine, as well as enzymes such as hydrolases or peroxidases, or the various chemiluminescers such as luciferin, or fluorescent compounds like fluorescein and its derivatives. In addition, the probes can be made inherently fluorescent as described in International Application No. WO 93/16094.

[00052] Various degrees of stringency of hybridization can be employed. The more stringent the conditions, the greater the complementarity that is required for duplex formation. Stringency can be controlled by temperature, probe concentration, probe length, ionic strength, time, and the like. Preferably, hybridization is conducted under moderate to high stringency conditions by techniques well known in the art, as described, for example, in Keller, G.H., M.M. Manak (1987) *DNA Probes*, Stockton Press, New York, NY., pp. 169-170.

[00053] As used herein "moderate to high stringency" conditions for hybridization refers to conditions that achieve the same, or about the same, degree of specificity of hybridization as the conditions "as described herein." Examples of moderate to high stringency conditions are provided herein. Specifically, hybridization of immobilized DNA on Southern blots with  $^{32}\text{P}$ -labeled gene-specific probes was performed using standard methods (Maniatis *et al.*). In general, hybridization and subsequent washes were carried out under moderate to high stringency conditions that

allowed for detection of target sequences with homology to sequences exemplified herein. For double-stranded DNA gene probes, hybridization was carried out overnight at 20-25° C below the melting temperature ( $T_m$ ) of the DNA hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. The melting temperature is described by the following formula from Beltz *et al.* (1983):

[00054]  $T_m = 81.5^{\circ}\text{C} + 16.6 \log[\text{Na}^+] + 0.41(\%G+C) - 0.61(\%\text{formamide}) - 600/\text{length of duplex in base pairs}.$

Washes are typically carried out as follows:

- (1) Twice at room temperature for 15 minutes in 1X SSPE, 0.1% SDS (low stringency wash).
- (2) Once at  $T_m - 20^{\circ}\text{C}$  for 15 minutes in 0.2X SSPE, 0.1% SDS (moderate stringency wash).

[00055] For oligonucleotide probes, hybridization was carried out overnight at 10-20° C below the melting temperature ( $T_m$ ) of the hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA.  $T_m$  for oligonucleotide probes was determined by the following formula from Suggs *et al.* (1981):

[00056]  $T_m (^{\circ}\text{C}) = 2(\text{number T/A base pairs}) + 4(\text{number G/C base pairs})$

[00057] Washes were typically carried out as follows:

- [00058] (1) Twice at room temperature for 15 minutes 1X SSPE, 0.1% SDS (low stringency wash).
- [00059] (2) Once at the hybridization temperature for 15 minutes in 1X SSPE, 0.1% SDS (moderate stringency wash).

[00060] In general, salt and/or temperature can be altered to change stringency. With a labeled DNA fragment of greater than about 70 or so bases in length, the following conditions can be used:

Low:	1 or 2X SSPE, room temperature
Low:	1 or 2X SSPE, 42° C
Moderate:	0.2X or 1X SSPE, 65° C
High:	0.1X SSPE, 65° C.

[00061] Duplex formation and stability depend on substantial complementarity between the two strands of a hybrid, and, as noted above, a certain degree of mismatch

can be tolerated. Therefore, polynucleotide sequences of the subject invention include mutations (both single and multiple), deletions, and insertions in the described sequences, and combinations thereof, wherein said mutations, insertions, and deletions permit formation of stable hybrids with a target polynucleotide of interest. Mutations, insertions, and deletions can be produced in a given polynucleotide sequence using standard methods known in the art. Other methods may become known in the future.

[00062] The mutational, insertional, and deletional variants of the polynucleotide sequences of the invention can be used in the same manner as the exemplified polynucleotide sequences so long as the variants have substantial sequence similarity with the original sequence. As used herein, substantial sequence similarity refers to the extent of nucleotide similarity that is sufficient to enable the variant polynucleotide to function in the same capacity as the original sequence. Preferably, this similarity is greater than 50%; more preferably, this similarity is greater than 75%; and most preferably, this similarity is greater than 90%. The degree of similarity needed for the variant to function in its intended capacity will depend upon the intended use of the sequence. It is well within the skill of a person trained in this art to make mutational, insertional, and deletional mutations that are designed to improve the function of the sequence or otherwise provide a methodological advantage.

[00063] PCR technology. Polymerase Chain Reaction (PCR) is a repetitive, enzymatic, primed synthesis of a nucleic acid sequence. This procedure is well known and commonly used by those skilled in this art (see U.S. Patent Nos. 4,683,195; 4,683,202; and 4,800,159; Saiki *et al.*, 1985). PCR is based on the enzymatic amplification of a DNA fragment of interest that is flanked by two oligonucleotide primers that hybridize to opposite strands of the target sequence. The primers are oriented with the 3' ends pointing towards each other. Repeated cycles of heat denaturation of the template, annealing of the primers to their complementary sequences, and extension of the annealed primers with a DNA polymerase result in the amplification of the segment defined by the 5' ends of the PCR primers. Since the extension product of each primer can serve as a template for the other primer, each cycle essentially doubles the amount of DNA fragment produced in the previous cycle. This results in the exponential accumulation of the specific target fragment, up to several million-fold in a

few hours. By using a thermostable DNA polymerase such as *Taq* polymerase, which is isolated from the thermophilic bacterium *Thermus aquaticus*, the amplification process can be completely automated. Other enzymes that can be used are known to those skilled in the art.

[00064] The polynucleotide sequences of the subject invention (and portions thereof such as conserved regions and portions that serve to distinguish these sequences from previously-known sequences) can be used as, and/or used in the design of, primers for PCR amplification. In performing PCR amplification, a certain degree of mismatch can be tolerated between primer and template. Therefore, mutations, deletions, and insertions (especially additions of nucleotides to the 5' end) of the exemplified polynucleotides can be used in this manner. Mutations, insertions and deletions can be produced in a given primer by methods known to an ordinarily skilled artisan.

[00065] The polynucleotide sequences of the instant invention may be "operably linked" to regulatory sequences such as promoters and enhancers. Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is "operably linked" to DNA encoding a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is "operably linked" to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is "operably linked" to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[00066] Polynucleotides and proteins. Polynucleotides of the subject invention can be defined according to several parameters. One characteristic is the biological activity of the protein products as identified herein. The proteins and genes of the subject invention can be further defined by their amino acid and nucleotide sequences. The sequences of the molecules can be defined in terms of homology to certain exemplified sequences as well as in terms of the ability to hybridize with, or be amplified by, certain

exemplified probes and primers. Additional primers and probes can readily be constructed by those skilled in the art such that alternate polynucleotide sequences encoding the same amino acid sequences can be used to identify and/or characterize additional genes. The proteins of the subject invention can also be identified based on their immunoreactivity with certain antibodies.

[00067] The polynucleotides and proteins of the subject invention include portions, fragments, variants, and mutants of the full-length sequences as well as fusions and chimerics, so long as the encoded protein retains the characteristic biological activity of the proteins identified herein. As used herein, the terms "variants" or "variations" of genes refer to nucleotide sequences that encode the same proteins or which encode equivalent proteins having equivalent biological activity. As used herein, the term "equivalent proteins" refers to proteins having the same or essentially the same biological activity as the exemplified proteins.

[00068] It will be apparent to a person skilled in this art that genes within the scope of the subject invention can be identified and obtained through several means. The specific genes exemplified herein may be obtained from root-knot nematodes. Genes, or portions or variants thereof, may also be artificially synthesized by, for example, a gene synthesizer.

[00069] Variations of genes may be readily constructed using standard techniques such as site-directed mutagenesis and other methods of making point mutations and by DNA shuffling, for example. In addition, gene and protein fragments can be made using commercially available exonucleases, endonucleases, and proteases according to standard procedures. For example, enzymes such as *Bal31* can be used to systematically cut off nucleotides from the ends of genes. In addition, genes that encode fragments may be obtained using a variety of restriction enzymes. Proteases may be used to directly obtain active fragments of these proteins. Of course, molecular techniques for cloning polynucleotides and producing gene constructs of interest are also well known in the art. *In vitro* evaluation techniques, such as MAXYGEN's "Molecular Breeding" can also be applied to practice the subject invention.

[00070] Other molecular techniques can also be applied using the teachings provided herein. For example, antibodies raised against proteins encoded by



polynucleotides disclosed herein can be used to identify and isolate proteins from a mixture of proteins. Specifically, antibodies may be raised to the portions of the proteins that are conserved and most distinct from other proteins. These antibodies can then be used to specifically identify equivalent proteins by immunoprecipitation, enzyme linked immunosorbent assay (ELISA), or Western blotting. Antibodies to proteins encoded by polynucleotides disclosed herein, or to equivalent proteins, can readily be prepared using standard procedures known in the art. The genes that encode these proteins can be obtained from various organisms.

[00071] Because of the redundancy of the genetic code, a variety of different DNA sequences can encode the amino acid sequences encoded by the polynucleotide sequences disclosed herein. It is well within the skill of a person trained in the art to create these alternative DNA sequences encoding proteins having the same, or essentially the same, amino acid sequence. These variant DNA sequences are within the scope of the subject invention. As used herein, reference to "essentially the same" sequence refers to sequences that have amino acid substitutions, deletions, additions, or insertions that do not materially affect biological activity. Fragments retaining the characteristic biological activity are also included in this definition.

[00072] A further method for identifying genes and polynucleotides (and the proteins encoded thereby) of the subject invention is through the use of oligonucleotide probes. Probes provide a rapid method for identifying genes of the subject invention. The nucleotide segments that are used as probes according to the invention can be synthesized using a DNA synthesizer and standard procedures.

[00073] The subject invention comprises variant or equivalent proteins (and nucleotide sequences coding for equivalent proteins or for inhibitors of the genes encoding such proteins) having the same or similar biological activity of inhibitors or proteins encoded by the exemplified polynucleotides. Equivalent proteins will have amino acid similarity with an exemplified protein (or peptide). The amino acid and/or nucleotide identity will typically be greater than 60%. Preferably, the identity will be greater than 75%. More preferably, the identity will be greater than 80%, and even more preferably greater than 90%. Most preferably, the identity will be greater than 95%. RNAi molecules will also have corresponding identities in these preferred ranges. These

identities are as determined using standard alignment techniques for determining amino acid and/or nucleotide identity. The identity/similarity will be highest in critical regions of the protein or gene including those regions that account for biological activity or that are involved in the determination of three-dimensional configuration that is ultimately responsible for the biological activity. In this regard, certain amino acid substitutions are acceptable and can be expected if these substitutions are in regions which are not critical to activity or are conservative amino acid substitutions which do not affect the three-dimensional configuration of the molecule. For example, amino acids may be placed in the following classes: non-polar, uncharged polar, basic, and acidic. Conservative substitutions whereby an amino acid of one class is replaced with another amino acid of the same type fall within the scope of the subject invention so long as the substitution does not materially alter the biological activity of the compound. Below is a list of examples of amino acids belonging to various classes

Class of Amino Acid	Examples of Amino Acids
Nonpolar	Ala, Val, Leu, Ile, Pro, Met, Phe, Trp
Uncharged Polar	Gly, Ser, Thr, Cys, Tyr, Asn, Gln
Acidic	Asp, Glu
Basic	Lys, Arg, His

[00074] In some instances, non-conservative substitutions can also be made. The critical factor is that these substitutions must not detract from the ability to manage nematode-caused diseases.

[00075] An "isolated" or "substantially pure" nucleic acid molecule or polynucleotide is a polynucleotide that is substantially separated from other polynucleotide sequences which naturally accompany a nucleic acid molecule. The term embraces a polynucleotide sequence which was removed from its naturally occurring environment by the hand of man. This includes recombinant or cloned DNA isolates,

chemically synthesized analogues and analogues biologically synthesized by heterologous systems. An "isolated" or "purified" protein, likewise, is a protein removed from its naturally occurring environment.

[00076] Recombinant hosts. The genes, antisense, and RNAi polynucleotides within the scope of the present invention can be introduced into a wide variety of microbial or plant hosts. Plant cells can be transformed (made recombinant) in this manner. Microbes, for example, can also be used in the application of RNAi molecules of the subject invention in view of the fact that microbes are a food source for nematodes

[00077] There are many methods for introducing a heterologous gene or polynucleotide into a host cell or cells under conditions that allow for stable maintenance and expression of the gene or polynucleotide. These methods are well known to those skilled in the art. Synthetic genes, such as, for example, those genes modified to enhance expression in a heterologous host (such as by preferred codon usage or by the use of adjoining, downstream, or upstream enhancers) that are functionally equivalent to the genes (and which encode equivalent proteins) can also be used to transform hosts. Methods for the production of synthetic genes are known in the art.

[00078] Where the gene or polynucleotide of interest is introduced via a suitable vector into a microbial host, and said host is applied to the environment in a living state, certain host microbes are preferred. Certain microorganism hosts are known to occupy the phytosphere, phylloplane, phyllosphere, rhizosphere, and/or rhizoplane of one or more crops of interest. These microorganisms can be selected so as to be capable of successfully competing in the particular environment (crop and other habitats) with the wild-type microorganisms, provide for stable maintenance and expression of the gene expressing a polypeptide of interest, and, desirably, provide for improved protection of the protein/peptide from environmental degradation and inactivation.

[00079] A large number of microorganisms is known to inhabit the phylloplane (the surface of the plant leaves) and/or the rhizosphere (the soil surrounding plant roots) of a wide variety of important crops. These microorganisms include bacteria, algae, and fungi. Of particular interest are microorganisms, such as bacteria, *e.g.*, genera *Pseudomonas*, *Erwinia*, *Serratia*, *Klebsiella*, *Xanthomonas*, *Streptomyces*, *Rhizobium*, *Rhodopseudomonas*, *Methylophilus*, *Agrobacterium*, *Acetobacter*, *Lactobacillus*,

*Arthrobacter*, *Azotobacter*, *Leuconostoc*, and *Alcaligenes*; fungi, particularly yeast, *e.g.*, genera *Saccharomyces*, *Cryptococcus*, *Kluyveromyces*, *Sporobolomyces*, *Rhodotorula*, and *Aureobasidium*. Of particular interest are the pigmented microorganisms.

[00080] Methods of the subject invention also include the transformation of plants or plant tissue with genes which encode the RNAi molecules of the present invention. In one embodiment, the transformed plant or plant tissue expresses antisense RNA and/or RNAi. Transformation of cells can be made by those skilled in the art using standard techniques. Materials necessary for these transformations are disclosed herein or are otherwise readily available to the skilled artisan.

[00081] Additional methods and formulations for control of pests. Control of nematode pests using the RNAi molecules of the instant invention can be accomplished by a variety of additional methods that would be apparent to those skilled in the art having the benefit of the subject disclosure. A "cocktail" of two or more RNAi molecules can be used to disrupt one or more of the genes identified herein. The "cocktail" of RNAi molecules may be specific to segments of a single gene or the entire gene. A "multigene cocktail" of RNAi molecules specific to two or more genes (or segments thereof) is also encompassed by the instant invention. In another embodiment of the instant invention, the disclosed RNAi molecules, cocktails, and/or multigene cocktails thereof, may be used in conjunction with other known nematode control agents and methodologies. Such cocktails can be used to combat the development of resistance by nematodes to a certain inhibitor or inhibitors.

[00082] Compositions of the subject invention which comprise RNAi molecules and carriers can be applied, themselves, directly or indirectly, to locations frequented by, or expected to be frequented by, nematodes. Microbial hosts which were transformed with polynucleotides that encode RNAi molecules, express said RNAi molecules, and which colonize roots (*e.g.*, *Pseudomonas*, *Bacillus*, and other genera) can be applied to the sites of the pest, where they will proliferate and be ingested. The result is control of the pest. Thus, methods of the subject invention include, for example, the application of recombinant microbes to the pests (or their locations). The recombinant microbes may also be transformed with more than one RNAi molecule thereby delivering a "cocktail" of RNAi molecules to the nematode pests. A carrier may be any substance suitable for

delivering the RNAi molecules to the nematode. Acceptable carriers are well known in the art and also are commercially available. For example, such acceptable carriers are described in E.W. Martin's *Remington's Pharmaceutical Science*, Mack Publishing Company, Easton, PA..

[00083] All patents, patent applications, provisional applications, and publications referred to or cited herein are incorporated by reference in their entirety to the extent they are not inconsistent with the explicit teachings of this specification.

[00084] Following are examples that illustrate procedures for practicing the invention. These examples should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted.

#### Example 1—Production of Hairy Roots for RNAi Testing

[00085] A hairy root assay system was developed for testing the anti-nematode activity of RNAi molecules.

[00086] *Agrobacterium rhizogenes*: Several *Agrobacterium rhizogenes* strains produce hairy roots on a variety of plant species. *A. rhizogenes* strains, A4, 15834, 8196 and LBA4404 demonstrate hairy root development on tomato and sugar beet, with A4 being the most efficient. The *A. rhizogenes* strain K599 demonstrated very efficient formation on transgenic soybean hairy roots and was also effective on sugar beet and *Arabidopsis*. However, stain K599 failed to produce hairy roots on tomato tissues possibly due to hyper-virulence.

[00087] Hairy root production: Transgenic hairy roots were identified by stable GUS expression in tomato, sugar beet, soybean and *Arabidopsis*. The construct pAKK1401 (pNOS / NPT-II / tNOS // pSU / GUS / tNOS) was used to produce hairy roots when transformed into *A. rhizogenes* strains A4 or K599. Transgenic roots were identified by GUS expression.

#### Example 2 – Protocol for Electro-competent *Agrobacterium* and Electroporation

[00088] Electro-competent *Agrobacterium* Protocol:

- [00089] 1. Grow *Agrobacterium* overnight in 5 mls LB + antibiotics at 30°C on shaker (for *Agrobacterium rhizogenes* strain K599 no antibiotics are needed).
- [00090] 2. Use the 5 mls of overnight culture to inoculate 500 mls LB + antibiotics at 30°C on shaker. Grow overnight.
- [00091] 3. Add liquid culture in eight 50 ml polypropylene orange cap tubes.
- [00092] 4. Centrifuge 10 min., 4000 rpm, 4°C.
- [00093] 5. Resuspend cells in each tube with 20 mls 10% glycerol (on ice)
- [00094] 6. Centrifuge 10 min., 4000 rpm, 4°C.
- [00095] 7. Resuspend cells in each tube with 10 mls 10% glycerol (on ice).
- [00096] 8. Centrifuge 10 min., 4000 rpm, 4°C.
- [00097] 9. Resuspend cells in each tube with 2 mls 10% glycerol (on ice).
- [00098] 10. Aliquot 50 µl into cold Eppendorf tube and place onto dry ice.
- [00099] 11. Store electro-competent cells at -80°C. These cells can be used for up to two years.

[000100] Electroporations:

- [000101] 1. Add 1 µl to 5 µl of DNA (resuspended in H<sub>2</sub>O and not TE or other buffer) to 50 µl of *Agrobacterium* electrocompetent cells and mix.
- [000102] 2. Transfer 20 µl of DNA/*Agrobacterium* mix to cuvette.
- [000103] 3. Electroporate:  
25µF, 400 Ω resistance, 2.5 volts (0.2cm cuvette) or 1.8 volts (0.1cm cuvette for BioRad electroporator. 330 µF, 4000 kΩ, low w, fast charge rate for BRL Electroporator.
- [000104] 4. Add 1ml of LB and transfer to Eppendorf tube.
- [000105] 5. Shake at 30°C for 2 hours.
- [000106] 6. Centrifuge down cells (2 min. 14 krpm).
- [000107] 7. Plate all onto LB + antibiotics (most *Agrobacterium* strains are naturally streptomycin resistant).

Example 3 – Protocol for Production of Transgenic Hairy Roots on Soybean

[000108] Seed Sterilization. Rinse the soybean seed with 70% ETOH for 2-5 min. Remove and add 20% Clorox and shake for 20-25 min. Rinse 3X with sterile water. Plate the seed, 5 seed per plate, onto  $\frac{1}{2}$  MSB5 + 2% sucrose + 0.2% gel (referred to as  $\frac{1}{2}$  MSB5). Place seed into chamber at 25°C, 16/8 photoperiod for 5-7 day (depending on genotype) germination period. After 1 week seedlings can be placed into cold room for longer storage if necessary (not to exceed 2 weeks).

[000109] Agrobacterium Preparation. For *Agrobacterium rhizogenes* strain K599, take a small sample from frozen glycerol into 25-50 ml of NZYM media with 50 mg/L kanamycin in a 125-250 ml Erlenmyer flask. Place onto shaker at 28-30 °C for 16 - 20 hours. Pour sample into centrifuge tube and centrifuge the bacterium at 4000 rpm for 10 min. Pour off supernatant and re-suspend the pellet with an equal volume of liquid  $\frac{1}{2}$  MSB5 + 200  $\mu$ M acetosyringone. Use pipette to re-suspend the pellet and homogenize the sample (remove all clumps). To determine O.D., prepare a 1:10 dilution by putting 900  $\mu$ l  $\frac{1}{2}$  MSB5 into cuvette and add 100  $\mu$ l of bacterial sample. Determine the O.D.<sub>660</sub> and calculate the volume needed to adjust (dilute) OD to approximately 0.2 for inoculation. Check final O.D.

[000110] Explant Preparation and inoculation. Place a sterile filter paper onto plates of  $\frac{1}{2}$  MSB5. Cut soybean cotyledons just above the shoot apex and place onto plate. Lightly scar the cotyledon's abaxial surface (flat side, upper surface that reaches toward sun) with a scalpel blade. Cut each cotyledon transversely into 2-3 pieces (no smaller than 1 cm). Add approximately 10 ml of prepared bacterial solution to each plate and allow cotyledons to incubate for 1 hr. Remove the bacteria using a vacuum aspirator fitted with sterile pipette tip, ensure that there is no standing liquid. Orient all explants with abaxial surface up and wrap plates for a 3 day co-culture, 25°C in light (16/8 photoperiod).

[000111] Hairy root selection and maintenance. After 3 day co-culture, wash explants with liquid  $\frac{1}{2}$  MSB5 + 500 mg/L carbenicillin. Transfer the explants abaxial side up to selection media,  $\frac{1}{2}$  MSB5 supplemented with 500 mg/L carbenicillin and 200 mg/L kanamycin. Roots should develop in approximately 2-3 weeks. The roots will form primarily from the cut vascular bundles with other roots developing from the small cuts on cotyledon surface. Remove roots (>1cm in length) and place onto replica media with

transfers to fresh media every 2 weeks to prevent *Agrobacterium* overgrowth. After 6-8 weeks on selection the roots can be moved to media without kanamycin, however carbenicillin must remain in media for several months for continued suppression of *Agrobacterium*. At this stage roots can be used for testing RNAi for nematode control. Sterilized nematodes can be added and observed for RNAi affects.

Example 4 – Testing of RNAi for Plant Parasitic Nematode Control.

[000112] Various types of nematodes can be used in appropriate bioassays. For example, *Caenorhabditis elegans*, a bacterial feeding nematode, and plant parasitic nematodes can be used for bioassay purposes. Examples of plant parasitic nematodes include a migratory endo-parasite, *Pratylenchus scribneri* (lesion), and two sedentary endo-parasites, *Meloidogyne javanica* (root-knot) and *Heterodera schachtii* (cyst).

[000113] *C. elegans*: RNAi vectors can be tested through expression of the RNAi in *E. coli*. *C. elegans* are fed *E. coli* and assayed for their growth by measuring growth of nematodes, production of eggs and viability of offspring. Another approach is to inject dsRNA directly into living nematodes. Finally, soaking nematodes in a solution of *in vitro*-prepared RNAi can quickly establish efficacy of treatment.

[000114] *P. scribneri*: The *P. scribneri in vitro* feeding assay uses a corn root exudate (CRE) as a feeding stimulus and both the red dye Amaranth or potassium, arsenate as feeding indicators. Feeding is confirmed after seven days by the presence of red stained intestinal cells in live worms exposed to the Amaranth or death of worms exposed to arsenate. This bioassay is used to test soluble toxins or RNAi. *P. scribneri* has also been cultured on wild type roots of corn, rice and *Arabidopsis*, and on A. rhizogenes-induced hairy roots of sugar beet and tomato. *P. scribneri* is very valuable in evaluating transgenic hairy roots because of the non-specific feeding of these worms.

[000115] *M. javanica*: Nematode eggs are sterilized using bleach and are used to inoculate hairy roots expressing RNAi. Nematodes are assessed for their growth by measuring knots, egg masses or production of viable eggs. An alternative approach is to microinject dsRNA directly into root feeding sites or into living female nematodes.

[000116] *H. schachtii*: Cultures of this nematode were maintained on sugar beets. Nematodes eggs are sterilized using bleach and used to inoculate hairy roots



expressing RNAi. Nematodes can be assessed for their growth by measuring knots, egg masses or production of viable eggs.

#### Example 5 – Plant Expression Vectors for RNAi

[000117] Modular Binary Construct System (MBCS): An important aspect of the subject disclosure is the Modular Binary Construct System. The MBCS eases the burden of construct development by creating modular pieces of DNA that can be easily added, removed, or replaced with the use of low frequency cutting restriction enzymes (8-base cutters). These constructs are useful for delivery of a variety of genes to plant cells and is not limited to the delivery of RNAi genes. To develop this system, a series of six, 8-base cutter restriction enzyme sites was placed between the left and right Ti borders of a previously created  $\text{kan}^R/\text{tet}^R$  binary plasmid (Figure 1). The production of both  $\text{kan}^R$  and  $\text{tet}^R$  MCBS aids the testing of constructs using different strains of *Agrobacterium rhizogenes* in different plant species. In addition to the MBCS, a series of shuttle vectors were created that aid in the cloning of useful DNA fragments by containing the multi-cloning site (MCS) of a modified Bluescript plasmid flanked by 8-base restriction sites (Figure 2). With six 8-base cutter sites, each site is, preferably, reserved for a particular function (Figures 3 and 4). Because of the close proximity of the *Pme* I and *Sgf*I sites to the left and right border of the binary vector, these sites are, preferably, reserved for gene tagging and enhancer trap experiments. The *Not* I site is, preferably, reserved for plant selectable markers (Figure 5). The *Pac* I site is reserved, preferably, for Plant Scorable Markers (Figure 6). The *Asc* I site is, preferably, reserved for RNAi experiments (Figures 7 and 8), while the *Sbf*I site is, preferably, reserved for anti-nematode proteins. The restriction sites that are denoted in the Figures are, preferably, reserved for the denoted insertions; however, the MCBS binary and shuttle vectors do not require the restriction sites to contain these suggested inserts.

[000118] Plant Selectable Markers for MBCS: To further develop the MBCS, a series of plant selectable markers were added to the MBCS (Figure 5). Plant selectable markers that were added to the MBCS include: pNOS/NPT-II/tNOS ( $\text{kan}^R$ ), pNOS/Bar/tNOS ( $\text{basta}^R$  for dicots), pUBI/Intron-Bar/tNOS ( $\text{basta}^R$  for monocots), and pUBI/Intron-PMI/tNOS (mannitol isomerase $^R$ ).

[000119] Reporter Genes for MBCS: Four exemplary reporter genes are used in the MBCS are provided in Figure 6 and Appendix 2. GUS, a nuclear localized GUS, GEP, and the anthocyanin transcriptional activator *papIC* genes into the MBCS.

[000120] Promoters for MBCS: We cloned several useful constitutive and nematode-inducible promoters (Figures 6, 7 and Appendix 2). Constitutive promoters include the SuperUbiquitin promoter from pine (pSU) and two promoter regions from the Strawberry Banding Vein virus (pSBV<sub>1</sub> and pSBV<sub>2</sub>). Seven nematode-inducible promoters from *Arabidopsis* were also been cloned.

[000121] The following Scorable marker clones have been constructed and placed in the MBCS, NPT-II binary vector (pNOS/NPT-II/tNOS):

Intron/GUS/tNos	Intron/NLS-GUS/tNOS	Intron/GFP/tNOS
pSU/Intron/GUS/tNOS	pSU/Intron/NLS-GUS/tNOS	pSU/Intron/GFP/tNOS
pSBV <sub>1</sub> /Intron/GUS/tNOS	pSBV <sub>1</sub> /Intron/NLS-GUS/tNOS	pSBV <sub>1</sub> /Intron/GFP/tNOS
pSBV <sub>2</sub> /Intron/GUS/tNOS	pSBV <sub>2</sub> /Intron/NLS-GUS/tNOS	pSBV <sub>2</sub> /Intron/GFP/tNOS
pKT/Intron/GFP/tNOS		
pKA/Intron/GFP/tNOS		

#### Example 6 — Control of Plant parasitic nematodes using RNAi in planta

[000122] Production of RNAi Vector. The RNAi shuttle vector to be used is adapted from the Modular Binary Construct System (MBCS - See Example 5). RNAi shuttle vectors preferably comprise a promoter, intron, antisense RNAi, stuffer fragment, sense RNAi, and terminator (See Figures 7 and 8 and Appendix 2 for more details). The plant promoter can be constitutive, tissue-specific or nematode-inducible. The intron is necessary to eliminate expression in *Agrobacterium*.

[000123] The anti-sense and sense RNAi molecules comprise nematode-specific sequences and are disclosed herein. These genes are associated with pathogenesis, growth, or other cellular function in nematodes. An exemplary group of RNAi sequences for use in plant/nematode control may be based upon:

[000124] 1. Genes specific for nematode esophageal gland cells.

[000125] 2. Genes specific for plant parasitic nematodes but not other free living nematodes.

- [000126] 3. Genes common to all plant parasitic nematodes.
- [000127] 4. Genes common to all nematodes (nematode-specific).
- [000128] 5. Genes specific for important tissues or cell types.
- [000129] 6. Genes from large gene families.
- [000130] 7. Genes involved in nematode signal transduction or other cellular pathways.

[000131] Appropriate RNAi constructs allow for the formation of dsRNA molecules (the sense and antisense strands join to form the dsRNA). The terminator sequence adds a poly-A tail for transcriptional termination. The RNAi shuttle vector can then be subcloned into the MBCS and transformed into *Agrobacterium rhizogenes*.

[000132] Plant Transformation with RNAi Vectors. An exemplary transformation system for generating hairy roots using *Agrobacterium rhizogenes* is provided below. The RNAi vector once introduced into the MBCS can subsequently (as a binary vector) be transformed in *A. rhizogenes* using, for example, the electroporation protocol of Example 2. Once the *A. rhizogenes* is confirmed to contain the plasmid, it is then used in generating hairy roots (See Example 3). Using this protocol transgenic hairy roots expressing RNAi are isolated, cultured and tested.

[000133] Testing of RNAi Vector for Nematode or Plant Pathogen Resistance. RNAi expressing hairy roots can be inoculated with sterilized nematodes. Infested hairy roots can be observed and the effect on nematodes determined. An alternative approach involves the microinjection of RNAi directly into root feeding sites (giant-cells for root-knot nematode, and syncytia for cyst nematodes) or into living female nematodes.

#### Example 7 – Insertion of Genes Into Plants

[000134] One aspect of the subject invention is the transformation of plants with genes encoding proteins of the present invention. Transformation of plants as described herein can be used to improve the resistance of these plants to attack by the target pest.

[000135] Genes, polynucleotides, and/or RNAi molecules as disclosed or suggested herein can be inserted into plant cells using a variety of techniques which are

well known in the art. For example, a large number of cloning vectors, for example, pBR322, pUC series, M13mp series, pACYC184, pMON, *etc.*, are available for preparation for the insertion of foreign genes into higher plants via injection, biolistics (microparticle bombardment), *Agrobacterium tumefaciens*, or *Agrobacterium rhizogenes*-mediated transformation, or electroporation as well as other possible methods. Once the inserted DNA has been integrated into the genome, the genetically modified-cell(s) can be screened via a vector carried-selectable marker that confers on the transformed plant cells resistance to a biocide or an antibiotic, such as kanamycin, G418, bleomycin, hygromycin, chloramphenicol, or bialophos, *inter alia*. The transformed cell will be regenerated into a morphologically normal plant. The transgene(s) in the transgenic plant is relatively stable and can be inherited by progeny plants.

[000136] If a transformation event involves a germ line cell, then the inserted DNA an corresponding phenotypic trait(s) will be transmitted to progeny plants. Such plants can be grown in the normal manner and crossed with plants that have the same transformed hereditary factors or other hereditary factors. The resulting hybrid individuals have the corresponding phenotypic properties.

[000137] It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application.

We claim:

1. An RNAi molecule, optionally comprising a linker, wherein at least one strand of said RNAi is encoded by a DNA sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 139.

2. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 1.

3. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 2.

4. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 3.

5. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 4.

6. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 5.

7. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 6.

8. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 7.

9. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 8.

10. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 9.

10. 11. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
12.
11. 12. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
12.
12. 13. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
12.
13. 14. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
13.
14. 15. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
14.
15. 16. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
15.
16. 17. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
16.
17. 18. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
17.
18. 19. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
18.
19. 20. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
19.
20. 21. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
20.

21. 22. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
22. 23. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
23. 24. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
24. 25. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
25. 26. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
26. 27. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
27. 28. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
28. 29. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
29. 30. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
30. 31. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
31. 32. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

33. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
32.
34. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
33.
35. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
34.
36. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
35.
37. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
36.
38. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
37.
39. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
38.
40. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
39.
41. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
40.
42. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
41.
43. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
42.



43. 44. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
44. 45. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
45. 46. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
46. 47. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
47. 48. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
48. 49. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
49. 50. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
50. 51. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
51. 52. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
52. 53. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
53. 54. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

54. 55. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
54.
55. 56. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
55.
56. 57. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
56.
57. 58. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
57.
58. 59. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
58.
59. 60. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
59.
60. 61. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
60.
61. 62. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
61.
62. 63. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
62.
63. 64. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
63.
64. 65. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
64.

65. 66. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
65.
66. 67. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
66.
67. 68. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
67.
68. 69. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
68.
69. 70. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
69.
70. 71. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
70.
71. 72. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
71.
72. 73. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
72.
73. 74. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
73.
74. 75. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
74.
75. 76. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
75.

76. 77. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
76.
77. 78. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
77.
78. 79. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
78.
79. 80. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
79.
80. 81. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
80.
81. 82. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
81.
82. 83. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
82.
83. 84. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
83.
84. 85. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
84.
85. 86. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
85.
86. 87. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
86.

88. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
87.
89. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
88.
90. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
89.
91. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
90.
92. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
91.
93. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
92.
94. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
93.
95. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
94.
96. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
95.
97. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
96.
98. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:  
97.

99. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 98.
100. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 99.
101. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 100.
102. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 101.
103. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 102.
104. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 103.
105. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 104.
106. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 105.
107. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 106.
108. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 107.
109. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 108.

110. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 109.

111. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 110.

112. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 111.

113. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 112.

114. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 113.

115. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 114.

116. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 115.

117. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 116.

118. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 117.

119. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 118.

120. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 119.

121. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 120.

122. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 121.

123. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 122.

124. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 123.

125. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 124.

126. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 125.

127. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 126.

128. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 127.

129. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 128.

130. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 129.

131. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 130.



132. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 131.

133. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 132.

134. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 133.

135. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 134.

136. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 135.

137. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 136.

138. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 137.

139. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 138.

140. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 139.

141. A transgenic plant or transgenic plant tissue comprising an RNAi molecule according to any of the preceding claims.

142. A method of disrupting cellular processes in a nematode comprising the steps of:
- (a) providing a composition comprising a compound according to any of the preceding claims; and
  - (b) contacting a nematode with said composition.

143. An isolated promoter comprising the following nucleotide sequence:

```
aacagcccaagataacagaaaagtcaaaggtgttcgaaa
gaccacttgtgactaaggatcatttcacccataattatctggtagca
cagactcatgataactgcgaggaacacaagttctttacagtcgattc
aaagacactttctctttacggtttcattgaaggagccgaccagaat
atgtcagagaagcttttctactgtgggttaatttcattaatctatcca
ggtgaaaacctcaaggagatctctcttctccaaaagacctctacag
ggcaatcaaaaactacagaaccagagtttgtagtgcacagagtagac
caatctacctgagaatcacgagtaccttcctagagtgggaaaatgat
gacatccttattccataccactggattgaggtaggactatccaatgg
aaaaattccatgggacaagtcataataagaagaccgcaacagtcgagt
atcttccagagataactgcactcagacctaaaaggataaaaagcagta
tataatcagtggtactaagatcttcgcagattcaaagaagaagcttaa
ctatgctgatgacaagataattctaataagcaattattcagaattaa
tcaaggagaaagaattaataactctttcagaatatgaagcccgcttt
acaagtggccagctagctatcactgaaaagacagcaagacaatggtg
tctcgatgcaccagaaccacatctttgcagcagatgtgaagcagcca
gagtgggtccacaagacgcactcagaaaaggcatcttctaccgacaca
gaaaaagacaaccacagctcatcatccaacatgtagactgtcgttat
gcgtcggctgaagataagactgaccccaggccagcactaaagaagaa
ataatgcaagtggtcctagctccacttttagctttaataattatgttt
cattattattctctgcttttgctctctatataaagagcttgatattt
catttgaaaggcagaggcgaacacacacagaacctccctgcttaca
aaccatgtattgtagctaaacctcttaggag.
```

144. An isolated promoter comprising the following nucleotide sequence:

tggtggggacaatggatccggtctgcgtagcaacaaggctg  
aaaaagattaaacagaaacctgtgatcattagcgttgaccaccacc  
aaaacctcctgagccaccaaaagcctccagagcctgaaaaaccaaagc  
ctccaccagcacctgaaccaccaaaagcatgtatgcaagccacctac  
tgcaacagttgtgatgttgtgtctgttactacctatgaaagtggag  
cggctgcaccattctttgagtcataatcgcgtagccatagccttcat  
gttaagtcctgtatttagccaataactaattcatcatgttctcatgct  
tttttgtttatttctttttctcaaatatgaatctctgttgtttgtcc  
ctccccctgtttataattagtcgcttctttgacacaagaagtctcatg  
agttcatgctaaagaaaataaaagttcaaattaaaacaccaaagtgt  
tgattaatctccataaacctgtgaagcagaaagttagtcattgtgac  
ctgaacagagcttaggaagtccttgaaggacatatcttcaagtgcta  
ttgggtcgtagcactcttaggcccatttaacttcattgagcccattaa  
attatgcaaaacaagaatgagacatatggaaacattagggttctta  
caggaaaaaataggaaaaagcagggacaactaaacaaaaattcagaa  
acaagaggcaagtggacgaccacggcgtgaagatcaacatgtggtgat  
gtgcatgagaccaagaccattttttctcggtcttcaacgcacacttg  
gtcttttcttatgtttgttgcatttctttatttaggcagaccctctct  
cttttttaattaggtatgtaaaaaatatgattttattttgttgaaa  
cattttgaggttaaaacctaaacttatagtaagcattttagtagagtga  
tttcttatacgacatctatcaacatgacctctaaccccccaaatatt  
gatgaaactactttaagtagtaaaacctaaagcaattaaaaatttctt  
ttaaatttagtagtttgtgtgtaatttaattgacatgattgcgtcgaaag  
aatcaaaacagttatatcgtgaacttaggagaatgttttatatcgt  
gtttcaacacatgattgctagcatatgtgtaggtgtcgtagacgtta  
cataacaatcatcactcgtaaatatcaaagtggtttctgagagaaac  
aaagggttatgattttcccaactgcactagttgtgtattgtttcttt  
cacacgtatgcttctgagttctgcccagaagtggaaattaaagcagag  
ttgggagagatcataatttattaggggtcgttatgctcaagtcatga  
cgtaaaatgaaaatttgtttttatttctttcaccaacacaaagaatag  
ctagttatctcttttttatatataacaattcatgaagttgatcagc  
tttatcacatcatccaatcgaattgctaatttagagatggaaatat  
caggatagagccaataagatatcaaattccaatggaccattttctcc  
atgtgctaattcatacaatctgtttttgtctgctttatttgatgatg  
atgctgagcgttttttaagtgtgaactaagatctagctaaccaaaaca  
aagatgggtctcttctgtctttgtcgtataagagcaagagagtggttt  
gattcaatttttaaaattctaaataaaaactccaaccgtgaatccagc  
catgaaactctttttagaaaatccttttttataacaaataattctct  
tgcttcttcttcttcttctgtttatttcaaccttttttggtttctttag  
ctcagaaaaagcccatctttttttctattcttgtttattttaatca  
tactgtgcgtttctacaaagtttgttcctttcttcttcaactctctc  
actcacagtcacagagatctgtttctttttctttttttgttttcaactc  
ttctcttccagt.

145. An isolated promoter comprising the following nucleotide sequence:

.agcaaagcaagaacaccagagaagaagaaaagcactacaga  
gaaaaatgtgagcttaagcgtctccaacaacacttctctgggagtc  
taaaggatgctgcaaaaagccttggtggtgagacttccgcataatttc  
caagcatgggtttatTTTTgttagcacacaaactatctgacctcga  
cttggattttctctcgcagtttgtccaactacattgaaacggatatg  
caggcaacatgggatcatgagggtggccatctcgtaagattaacaaag  
tgaacagggtcactaaggaaaatacagacgggtactggactcgggtcaa  
ggtgtagaaggaggactaaagttcgactcagcaactggcgaattcat  
tgcagttagaccttttattcaagaaattgatacccaaaagggctctgt  
cgtctcttgataatgatgcacatgcaagaagaagtccaggaggatatg  
cctgacgatacttcattcaagctccaggaagctaaatctgtcgacaa  
tgccattaagtttagaggaggatacaaccatgaatcaagcaagaccag  
gtaagaacttctctatccataaaccatagatggagcgaattagaatct  
  
taatccattttcagtttttgcaggatcattcatggagggttaatgcta  
gtggtcagccatgggcttggtatggccaaagagtctggcttgaatggc  
agtgaaggaataaagagcgtttgcaacttaagctctgtggaaatttc  
agatggaatggatccaacaatccgatgcagtggcagttattgttgaa  
ctaaccaatccatgtcatgcagcatatcagattcatcaaatgggtca  
ggcgcagttctgcgtggaagctcatctacttccatggaagattggaa  
ccaaatgagaaccacacaacagtaatagcagcgcagagtggatcaacaa  
cgctgatcgtaaggccagttatagagaagacactgtacgtttcaag  
ttcgagccatcagttgggtgtcctcagctctacaaagaagtggaaa  
acgttttaactgcaggacgggtcgtttcagctgaagtacttggatg  
atgaagaagaatgggtgatgctggttacagattctgatctccaagaa  
tgtttggagatattacatggtatgggaaaacactcgggtgaagtttct  
cgttcgtgatttgtctgccccctctaggtagttctggtggcagtaatg  
gttatcttggaaacaggcttatgacgtcgtaagacatagacacacaca  
gttatgtattcccagtgaaagaatgttggttatttctctagatatta  
gtatgcttataaataggcatgaaggagaaagacaattttggtatagt  
ggagttcagcagaaaaatgtatatgtttttcgttttatatgaatcag  
agaataaaaagttggatgttatatctacgttgctaagtgtgtacctgc  
tcacccatctttcatataagaaaagagaacacttttagttatccctg  
tgatgcagaatcgtatttcttggttatctctccattcctgtggaaacc  
aaciaagtcaactaaatttcggtttaattgggtgggtttttaagtcaa  
cgaggacttgatttttagttgggcttgggcctataattgtgttcatca  
ttgggttttttcccccttatcagtttaacgtccatatccatatcttt  
ttcttttttaacggcaaagttcatatccatatcttatgatgtgcct  
aaaagaggggagaagatgcgaagacagaattttcatatttgaaagggt  
tcgatatcgatattgggaaacgaatcaagggtcaaaaaactcagttca  
atagttgaaatttaaaaattttattaattcaatccgattgggttcgt  
tttggttatgggttcggttctatatcatcaaccaatcggtttgggtcct  
aaagataattataaatattcaccaacaccagtggttaaacacatatca  
acaaacctaaagttagataaacaagaga.

146. An isolated promoter comprising the following nucleotide sequence:

aaattggcactcttcttctgctgggttccaaaagaaacgaat  
caatatgtgcaacaagaagagctccagaagcagtcattctctaaat  
cttaatctaacaacagctcaagaagaaaaaattccatagctagaga  
gaacacaaagtcacaagacgacgtcgtagaggcacaagtcaaacct  
gaatggcttaagccgaactgagtggttttgactagaccatcatcaga  
aaagtctccaagacggtagtcggatgttagatcgctcaagtaatttt  
tgggttttggttggtctcacgttttcagctgcccatttgatttcagttt  
gggcttttccttatctctaaaggccaatttcatttaggtttagttt  
atgtgatcattatccttactataaaggcttcgcctttcgagaaattt  
agggtttcttctgtctgtctcgctcactcaggtttgtgcctcaacgac  
tgcttcacttctagcttgattcttcttcttcgtttatatgtatactg  
tacattagattattcttggtttctcgagcttctgctatagattttgat  
tcttttttttggtgtcttggtttcgtttccaggatcagatcttagct  
aaattgagacaagctcaaaatgaggtacttgacgcacatctcttaoatt  
cactgtttaattagagaacaatacgtctctgaatcgtgattcagaga  
cgtattgttcttctgtcatatgcaataagtttaattagagaacaata  
cgtctctgaatcgtgattgttttttggtgtgcgttattgatagctt  
tatgatgttaatagtctaggattgacacgaagtgttctgcagtttt  
gcataaatgctcttactaaggcctctaaatttggtgacaaatcta  
aatcttgccctataaaaaatttaggtgtattaagataagattattttg  
tatggtagtgtctataatgtgggttggtcatgttgaggttgatcaatg  
ttgtgtatttttggttggttagttaatttgcttaactctgttctttg  
tgggttaatacagtaagcttcagagtgaggccggttcgtgaagccatc  
actactatcacagggaatccgaggcaaaagaaacgtaactttgtcga  
gactattgagctccagatcggctctgaagaactatgaccctcaaaagg  
acaagcgtttcagtggtatctgtcaagttaccacatatccccgcctc  
aaaatgaagatctgcatgctcggagatgccagcatgttgaagaggt  
gatatatctttcatggaaattgatcattttgtgctctgttcttctg  
ataatgggttttggtgctcatttcatttgggtggctctattagtttcatt  
tgatgttgatataatgtcttctgaatgtagatgcatgatgttttcggaa  
tttgggtcattgtttatttaggcttcatttcttgcataattaaatatt  
tgcttatttcatcttgatcttttcgtaggctgagaagatggggttg  
gaaaacatggatgttgagtctctaaaaaagcttaacaagaacaagaa  
actcgtcaagaagcttgcaaagaaataccatgctttcttggcctctg  
agtctgtcattaagcagattcctcgtcttcttgggtcctggctttaac  
aaggcaggcaagttctggctacagctaataattccattgttcttcttt  
acatccgttttgatgtttggataggttttagtagtctatttcttttgt  
caatgtctttttgatataatgccaatcctttatcctgtgagattatg  
cttctttgatgattcttaagtaacattcctttgctttactttacaca  
ggaaaattcccaactcttgtagccaccaggaatccttggagtcaaa  
gggtgaatgaaacaaaggcaacagtgaaagttccagctgaagaaggttc  
tgtgcatgggagttgcagttggttaacctt.

147. An isolated promoter comprising the following nucleotide sequence:

```
ttggcaaactgagatataagaggggaaggtgat tttcatgcaa
at tttttttttttttttttttttgaatgaatgcaaaatttat tcaaaaa
aaaaaaacctgggctacatcaagtacttcatttctgagtttttgaaa
aatctaaagacaacaaaagactttacaatttaataaaaaaataataa
aaatactttatcactctcaacgaaattgttgatttaataacgtatct
cttggtaaaacagcgtttttatttgacgaaattgttataaatgaataa
aatgataatagaaaactagtgtggtacgtaaaatacctctcatttggc
aaaataacggttatgtatcatgagatttgcatacgacagcgtgctta
aatagtgtgctttcaggagaaaaatatataccaagttatttgctgaaa
ttaccacgcaaactctgaggttcgaatggcaaaataaaaaaccaatgt
catttccttaattgtattaaggtcatttaataaaaattgtacactttt
ttcacctgtaagcgttccaaagtgtagaatggataactagaagggtc
aaaggtataatattaataagcgaactcactttttgcccagtgattt
cacttcttacattttgcttgatatagttacccaaaagtgtatatatat
tcccttatacaattgttctattttctggattataaggggaataagaa
aaaagaaaagagagagtataataataactttttataaagtgatgtta
gattctaat tttgtaacgaaaagttcaaagtgaaagaaaaaacgaaaa
agtttttctgtttttgttttatatctatagccaagaaagtttctcaga
tttacaagaagttaactgagaaaaacaaaaaaaaaacttatgaagca
tgaaagactaatttaacgaggtgatttaattttgagacaaattaaacat
cgaattaaaagtaacattttggaggggtttatatgttatatatgtgaca
tgataagtcgattcatgactaatgtatatctggaatctaactatgga
agaatagagaacgaagcagagccaaggtcaacttgccagacacgaat
caacagatttgtgaatgagaccaaataaatgggtcataaaccggttggg
tttaaacgggcaagtcattccttgggtcaattccattcgttattcctt
catgcaagaccctctgatacaaccaaagactcccattacaatatctt
ttcgatcacgagctacttattttcaaattgtgttacctctttcggtgac
tcttgtgttgtgtggtaaagcctagtcgagatgtgtcggtatatata
ggcatacatatacaaatgcgacaaaataagtatatatttatattgtttaa
tttctatatattccatttctatatgcatgggtgggatttttgaccaaaa
ccctaattcaagaatagaatccaaaagatgggatcaaagaatataat
ctaattgggctgaccacattttccgatttaattcgcatagttaatatt
ctttccactactttatgccgcagaaatttgtaattaaagtaagacaaa
gaaatacagatatagaatgggtcgtagaaaccagtagaggaatttcat
ttttcggtggataagtgggaatattaataagagaatgggtctttactctt
tacagtgggaaatgggaatagtagccattataatttcatcagattc
tatatatgcatgtttgtataagctaaaataaaatacgtttaagcatc
ttcaaaaaaatttacaagttctagagactctcttaacgtcggcaatt
tatattctactttacatgacactttcaggaaaagaaaactataactca
ctagcagatcattaaattttctttttctttttttgaatgaaccttag
ttgtgggtttttattttttgttagctagaaaacttcagtggtttttttcc
gccaatggtagtgctttgatgatgggtccgg .
```

148. An isolated promoter comprising the following nucleotide sequence:

caatcaaggtaacgaaggaggatcagcgaaaggatgggcta  
tat ttggagtttttctcgtgtaagtaatgctttgtgatcttcca  
tgccgacatataactgaagaataaactcaactcattgtgttctggtg  
tgtttcttctgatcagattcctcgttgcatctgcacttttctgctgt  
gggggctttatttataaaacaagagtagagcgtgtggtaatcttcat  
atctttctacaattccacttccattctctaattattctctcacgtga  
tatacacacactcaatcactgatgtactcgtatggatgcagcgtgga  
actgatgcattgccggggatgtcacttctatcgggcttactagaaac  
tgtaagtattacaagaaaactcaaaaggattccatttatgcaaatc  
taagagaaagctcactgtggtctttggttacaatttatggatctctc  
aagagacaaatgctatgtaagctaattgatttttggcttgataaaca  
ggtagagtgaagtggacaaagctactcaagaactgaagacatcaaca  
atgcttttgccaatgaagtctcatgggaccgctcttccgcatcttct  
actcaagcgacaacaacacagagaccaagtgaagaacatatggtgc  
gatctaattttgtcaagtgcctcacaagaggtagctttcaagccat  
ggtatggcacgcttgtgatctgcgatttctggattttgctttgtatg  
tttattttctaccttctagaaagagggtcaaaaagttaatagcttcac  
cgtgagaatggtgttttaccagattcatgtgctatgatagaaaaag  
acaaagcaaacaagagttctttctttgcttaggttacaagaacaaga  
gtatcgttataaagtcaacaaagattgaaacatattttgtcaaggg  
agtgggttagaatctcttctactctcttgcccttctcactaagacaa  
aaaaaagacttggactttgtctaagggttttgtggatattattaacca  
agtccttttgcaaaaagtaatatgttttttcgcattcctcttttag  
aatttagtttaattctaggcttttatattgggttattactttcttgaaaa  
atgatctgtttattctattcactacttggttacctcgctttttatctt  
acttctacaaaaggattatcagtgaaagttagtctcttactctcacc  
ttccgaaaaataaaacaaaaatatcgatacttctagatcaaaccaagt  
tgattaaaacatccctattccctacgattctgatcttgagatatatt  
atcatgttaagatctaaattgacaagaaaactgatttttcatctcta  
gtaggaaaaataattactattagtgatcatgattgtcgaccgtaaga  
ggtgggttagttactctccatctttctttgaagaagtcagaaagtc  
gaaattatatcaaattaaacatcaatattgaacacatatatctgtat  
ggttttatgtttagaaaattccaatatttatatattcctagggaaaa  
agaagcttattcttcaaattattgttatgagtcggtaaaatatggat  
aaaaatataaagtctaaatatataaaactcagtttgctttgctttta  
cctctccaagtcctccaaagtc aaattatttttagttaattaaaccaa  
aaaagggtttattagtc aaacttagcatgcaatgctgggtaccaaacc  
caagcatttagtctcttttaattctctttttctccaataagtttttac  
aatttttaattgtttgcaattcccttgattatttatcttcatcccaa  
tttagctaataccaactccgtttcttattcttccaagtcttttctta  
taaatacgttcttcttccctcttatttcatatcactcaccacaaag  
tcttctcatttctcat .





150. An isolated promoter comprising the following nucleotide sequence:

gattaggggtttgagttgtcactggaagaggtttgattgt  
gagtgatgatggagagattatgaaggagtttgtgtgtatttatagag  
gagttaggggttttgaggtttgatgagaagtaggtttgaagaagtttt  
gttgttgcaacttatttagagttacttgttccacaaccacaagtaag  
attgggtcacttctaagttctaactagaaacaacccatgacacatggag  
atctcagctaacctagtttaattgtatatgtattatattttatttaa  
tattataaaaataaaaataaattttcacaaataaaagaactacaaaaa  
gtgagaaaaataatttgataaacaatttagaaaattagtatatcaa  
taaaataaatttataatccgatgggttttgccctttgggtttggcctttg  
tttgaacttcgatgagtgactatgtatagcgaaaacaattcgggttg  
tttttggtttaatttttaaaaaatacaagcgacaatatctgatgagaa  
taggtgaaaagcaaataatatcagtttaattggaaatatttactttt  
ttacaactaatattttgtttgggtcaaccaacaaatagatttaattaa  
ttatgggtttatgagcttttattttgttgcgacagtatatatatgttaa  
aatagtgatattgcatggcggaagggtccggaagcaacacatatctcc  
tttttaatttttttttaacaagaataacatgttaatttttttttgga  
aattaataaagaatacatatttctaatttttgcgtcagatagatgat  
taaagagtgtgtgttttttttaacaacaaggaatacattatacata  
tttcataatttctctcgacattgtttgttttttttaaaaaatagattaa  
agagtcctacgaagctaagtagctaacgaagacttgaaatgagaagaa  
gacgagaatcttttaatatatttttggttaagcgataatatttgaaaa  
ttaataaatatagatttaaggaaataacaataacgcagatatcggtaa  
gtcatagaaaaaaagaaacaacacaaacttacataaacatgtttcct  
aatttgtaatggagtaaaattccttcttttttttttttttttgattt  
ggattccaattagtaaagaactcaatgactataaataacctttaacc  
ctctcattatttcttactatcaattgattaagctctcgttcctaaga  
aagcaatagacgaacaagaacccatcgaagaacacaaatctctcttt  
gaagttgtcgataatgttagtacaccgttacttcgtccaagactttt  
ttgccgttcggtttcttacaaaacaaggatttggttaccattacttt  
tgtcgtaactcctttttacatgtacgtcaaaaagtgggttcctcgctc  
cggccttgagaaacgacacctcttaccacaaaaagcttattttaaac  
cgtctaaaaccggaaaatctcaatctaaaccggatcgggttcattgag  
aaaccgattcaaacaccgagtgagaagtagaattttttgatgggtc  
cgtcacaatgtgtgtgctgctccttcgcccaagacatgtaccgattccga  
tattttgtgggtgtaaagatgatcaaagagtcctcaaagctaagcacg  
acttgaatgagaagaagaagaccaattactcaattagattttgtttt  
gtggagcaattattgtctatttatctttgttttttagcaaataatctg  
tatccactaatcttcacagtacttgactaacaagaagtaaagagttt  
tcttattttccaattgttttttaattctgatacttttttcataatttta  
caatgtttgatgaaaaaaaacattcaaaccctaaattttctttttttg  
gtatgaattcaaaccctgaattacttttgacgaggaccgcggtata  
aatagggtgatctcccaacaacaaaaaggt.

151. A transgenic plant or transgenic plant tissue comprising an isolated promoter according to any of claims 143 through 150.

54  
**APPENDIX 1**

SEQ ID NO:	INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTIDE / GENE
1, 2, 3	2293133	glyceraldehyde-3-phosphate-dehydrogenase
4, 5, 6, 7	7143495	Histone H4
8 & 9	7143515	ATP dependent RNA helicase, mRNA sequence
10, 11, 12, 13	7143527	nematode specific
14 & 15	7143602	protein serine-threonine phosphatase 1, catalytic subunit
16 & 17	7143612	40S ribosomal protein S4
18	7143666	cytochrome p450
19, 20, 21, 22	7143675	Neuroendocrine protein 7B2
23, 24, 25	7143839	nematode specific
26	7143863	40S ribosomal protein S17
27 & 28	7144016	vacuolar ATP synthase subunit G
29	7144025	malate dehydrogenase
30 & 31	7144060	J2 pcDNAII Globodera rostochiensis cDNA similar to Bystin, mRNA sequence
32 & 33	7144225	similar to arginine kinase
34	7144354	pyrroline-5-carboxylate reductase

SEQ ID NO:	APPENDIX 1 (cont.) INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTIDE / GENE
35, 36, 37, 38	C10	ribosomal protein L18a
39, 40, 41, 42, 43	C118	ribosomal protein S11
44 & 45	C122	ribosomal protein L16/L10E
46 & 47	C127	FMRFamide-related neuropeptide precursor
48	C129	ADP-ribosylation factor 1
49	C130	ribosomal protein L11
50	C137	nematode specific; conserved in <i>C.elegans</i>
51 & 52	C138	ribosomal protein L7
53	C145	ADP/ATP translocase
54 & 55	C148	troponin
56 & 57	C154	calponin
58	C16	translation elongation factor EF1A
59 & 60	C18	40S ribosomal protein S16
61	C27	ubiquitin
62 & 63	C46	nematode specific
64, 65, 66	C48	ribosomal protein S3AE
67	C59	40S ribosomal protein S5/S7

SEQ ID NO:	<u>APPENDIX 1 (cont.)</u> INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTIDE / GENE
68	C8	glyceraldehyde 3-phosphate dehydrogenase
69 & 70	C82	60S ribosomal protein L30/L7E
71	C90	glyceraldehyde 3-phosphate dehydrogenase
72	C135	nematode specific
73 & 74	C206	predicted troponin
75	C227	cytochrome P450
76	C238	vacuolar ATP synthase subunit G
77	C246	40S ribosomal protein S4
78	C308	FMRFamide-like neuropeptide precursor
79	C342	ubiquitin
80 & 81	C344	nematode specific; conserved in <i>C.elegans</i>
82, 83, 84, 85	C370	40S ribosomal protein S5/S7
86	C426	nematode specific
87	C458	histone H4
88 & 89	C481	ribosomal protein L30E
90 & 91	C556	nematode specific; conserved in <i>C.elegans</i>

SEQ ID NO:	APPENDIX 1 (cont.) INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTIDE / GENE
92	C628	ribosomal protein S17E
93 & 94	C665	malate dehydrogenase
95 & 96	C669	malate dehydrogenase
97	C694	ribosomal protein S3AE
98 & 99	C709	ADP/ATP translocase
100 & 101	C714	ADP-ribosylation factor 1
102	C721	calponin
103 & 104	C726	ribosomal protein L11
105	C736	nematode specific
106 & 107	C773	troponin
108	C834	nematode specific
109	C860	bystin
110 & 111	C863	troponin
112 & 113	C883	translation elongation factor eEF-1A
116	C888	40S ribosomal protein S16
117	C898	glyceraldehyde 3-phosphate dehydrogenase
118 & 119	C935	peptidyl-glycine alpha-amidating monooxygenase
120 & 121	C937	calponin
122 & 123	C942	peptidyl-glycine alpha-amidating monooxygenase

SEQ ID NO:	<u>APPENDIX 1 (cont.)</u> INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTIDE / GENE
124	C954	arginine kinase
125, 126, 127	C969	calponin
128 & 129	7235653	ribosomal protein L18A
130	8005381	neuroendocrine protein
131	7235496	pyrroline-5-carboxylate reductase
132 & 133	7275710	protein phosphatase pp1-beta catalytic subunit
134	7923685	nematode specific
135	7641370	40S ribosomal protein S11
136 & 137	7923404	nematode specific
138	7797811	ATP-dependent RNA helicase
139	7143613	predicted phospholipase D

## Appendix 2:

### Exemplary genes used for RNAi vectors.

#### Promoters:

##### *Constitutive:*

##### **Super Ubiquitin from Pine**

CCGGGGAAACCCCT CACAAATACATA AAAAAAATCTT TATTTAATTATC AAACCTCTCCACT ACCTT  
 TCCCACCAACCGTTA CAATCCTGAATG TTGGAAAAAACT AACTACATTGAT ATAAAAAACTA CATT  
 CTTCTPAAATCATAT CAAAATGTGATA AATATATCCACT CAAAGGAGTCTA GAAGATCCACTT GGACA  
 AATTGCCCATAGTTG GAAAGATGTTCA CCAAGTCAACAA GATTTCATG GAAAAATCCATC TACCA  
 AACCTACTTTCAAGA AAATCCAAGGAT TATAGAGTAAAA AATCTATGTATT ATTAAGTCAAAA AGAAA  
 ACCAAAGTGAACAAA TATTGATGTACA AGTTTGAGAGGA TAAGACATTGGA ATCGTCTAACCA GGAGG  
 CGGAGGAATTCCTTA GACAGTTAAAAG TGGCCGGAATCC CGGTAAAAAAGA TTAAAAATTTTT TGTAG  
 AGGGAGTGCTTGAAT CATGTTTTTAT GATGGAAATAGA TTCAGCACCATC AAAAAACATTGAG GACAC  
 CTAAAAATTTTGAAGT TTAACAAAAATA ACTTGGATCTAC AAAAATCCGTAT CGGATTTTCTCT AAATA  
 TAACFAGAATTTTCA TAACTTTCAAAG CAACTCCTCCCC TAACCGTAAAAAC TTTTCTACTTC ACCGT  
 TAATTACATTCCTTA AGAGTAGATAAA GAAATAAAGTAA ATAAAGTATTG ACAAAACCAACAA TTTAT  
 TTCCTTTTACTT AAAAAACAAAA AGTTTATTATT TACTTAAATGG CATAATGACATA TCGGA  
 GATCCCTCGAACGAG AATCTTTTATCT CCTGGTTTTGT ATTAAAAAGTAA TTTATTGTGGGG TCCAC  
 GCGGAGTTGGAATCC TACAGACGCGCT TTACATACGTCT CGAGAAGCGTGA CGGATGTGCGAC CGGAT  
 GACCTGTATAACCC ACCGACACAGCC AGCGCACAGTAT ACACGTGTCTATT TCTCTATTGGAA AATGT  
 CGTTGTATCCCGC TGGTACGCAACC ACCGATGGTGAC AGGTCTGTCTT GTCTGTCTGCGT AGCGG  
 GAGAAGGGTCTCATC CAACGCTATTAA ATACTCGCCTTC ACCGCGTTACTT CTCATCTTTTCT CTTCG  
 GTTGATAATCAGTG CGATATTCTCAG AGAGCTTTTCAT TCAACCCGGG

##### **Strawberry Banding Vein Virus 1**

aagcttttcactgtgggttaatttcattaatctatccaggtgaaaacctcaaggaga  
 tctctcttctcccaaagacctctacagggcaatcaaaaactacagaaccagagttt  
 gtagtgacagagtagaccaatctacctgagaatcacagtagccttcctagagtggtg  
 aaaatgatgacatccttattccataaccactggattgaggtaggactatccaatggaa  
 aaattccatgggacaagtcatataagaagaccgcaacagtcgagtatcttcagaga  
 taactgcactcagacctaaaaggataaaaagcagtatataatcagtgtagtaaatct  
 tcgcagattcaagaagaagctt

##### **Strawberry Banding Vein Virus 2**

Gtttaacaacagcccaagataacagaaaagtcaaagggttcgaaagaccacttgt  
 gactaaggatcatttcattccataattatctggtagcacagactcatgataactgcga  
 ggaacacaaagttctttacagtcgattcaaagacactttctctttacggtttcattga  
 aggagccgaccagaatattgtcagagaagcttttactgtgggttaatttcattaat  
 ctatccaggtgaaaacctcaaggagatctctcttctcccaaagacctctacagggc  
 aatcaaaaactacagaaccagagtttgttagtgacagagtagaccaatctacctgag  
 aatcacagtagccttcctagagtggtgaaatgatgacatccttattccataaccactg  
 gattgaggtaggactatccaatggaaaaattccatgggacaagtcatataagaagac  
 cgcaacagtcgagtatcttcagagataactgcactcagacctaaaaggataaaagc  
 agtatataatcagtgtagtaaatcttcgagattcaaagaagaagcttaactatgc  
 tgatgacaagataattctaataagcaattattcagaattaatcaaggagaaagaatt  
 aataactctttcagaatatgaagcccgctttacaagtgccagctagctatcactga  
 aaagacagcaagacaatggtgtctcgatgcaccagaaccacatctttgcagcagatg  
 tgaagcagccagagtggtccacaagacgcactcagaaaaggcatcttctaccgacac  
 agaaaaagacaaccacagctcatcatccaacatgtagactgtcgttatgcgtcggct  
 gaagataagactgacccagggccagcactaaagaagaataatgcaagtggctcctag  
 ctccacttagctttaataattatggttcattattattctctgcttttgctctctat  
 ataaagagcttgatattttcatttgaaggcagaggcgaacacacacacagaacctccc  
 tgcttacaaccatgtattgtagctaaacctcttaggaggatattc



**Nematode Inducible:****Trypsin Inhibitor from Arabidopsis (clone#6598343)**

cccgggagcaaagcaagaacaccagagaagaagaaaagcactacagagaaaaatgtg  
agcttaagcgctctccaacaacacttctctgggagctctaaaggatgctgcaaaaagc  
cttgggtggtagacttccgcatatctccaagcatgggtttatttttggtagcacaca  
aactatctgacctcgacttggattttcttctgcagtttgccaactacattgaaac  
ggatatgcaggcaacatgggatcatgaggtggccatctcgtaagattaacaaagtga  
acaggtcactaaggaaaatacagacgggtactggactcgggtccaaggtgtagaaggag  
gactaaagttcgactcagcaactggcgaattcattgcagtttagaccttttattcaag  
aaattgatacccaaaaggggtctgtcgtctcttgataatgatgcacatgcaagaagaa  
gtcaggaggatagcctgacgatacttcaatcaagctccaggaagctaaatctgtcg  
acaatgccattaaagttagaggagatacaacatgaatcaagcaagaccaggtaaga  
acttctctatccataaaccatagatggagcgaattagaatcttaatccattttcagtt  
tttgaggatcattcatggaggttaattgctagtgggtcagccatgggcttggatggcc  
aaagagctctggcttgaatggcagtggaaggaataaagagcgtttgcaacttaagctct  
gtggaaattttcagatggaatggatccaacaatccgatgcagtggcagttattgttgaa  
cctaaccaatccatgtcatgcagcatatcagattcatcaaattggctcaggcgcagtt  
ctgcgtggaagctcatctacttccatggaagattggaaccaaatagagaaccacaac  
agtaatagcagcgagagtggtatcaacaacgctgatcgtaaggccagttatagagaa  
gacactgtacgtttcaagttcgagccatcagttgggtgtcctcagctctacaaagaa  
gttggaacacgttttaactgcaggacgggtcgtttcagctgaagtacttggatgat  
gaagaagaatgggtgatgctggttacagattctgatctccaagaatgtttggagata  
ttacatgggtatgggaaaacactcgggtgaagtttctcgttcgtgatttgcctgcccct  
ctaggtagttctgggtggcagtaattggttatcttggaaacaggcttatgacgtcgtgaag  
acatagacacacacagttatgtattcccagtgaaagaatgttgtttatttctctaga  
tattagtagctttataaataaggcatgaaggagaaagacaattttggtagtgaggt  
tcagcagaaaatgtatatgtttttcgttttatatgaatcagagaataaaagtggga  
tggtatatctacgttgctaatgttgtacctgctcacccatcttcatataagaaaag  
agaacacttttagttatccctgtgatgcagaatcgtattctttgttatctctccatt  
cctgtggaaaccaacaaagtcaactaaatttcgggttaattgggtgggtttttaagtc  
aacgaggacttgatttttagttgggcttgggctataattgtgttcatcattgggttt  
ttcccccttatcagtttaacgtccatatccatatctttttcttttttaacggcaa  
agttcatatccatatcttatgtgtgctgctaaaagggagaaagatgcgaagacagaa  
ttttcatatttgaaaggttcgatatcgatattgggaaacgaatcaaggtcaaaaaa  
ctcagctcaatagttgaaatttaaaaaattttattaattcaatccgattgggttcggt  
ttgttatgggttcggttctatatcatcaaaccaatcgggttggtcctaagataatta  
taaatattaccaacaccagtggttaaacacatatcaacaaacctaaagttagataaa  
caaagagacccggg

**Arabidopsis Transmembrane Protein from Arabidopsis  
(clone#6468048)**

cccgggaattggcactcttcttctgctgggttccaaaagaaacgaatcaatatgtgc  
aacaagaagagctccagaagcagtcattctctaaaatcttaatctaacaacagctca  
agaagaaaaaattccatagctagagagaacacaaagtcaacagacgacgtcgtaga  
ggcacaaggtcaaacctgaatggcttaagccgaactgagtggttttgactagaccat  
catcagaaaaggtctccaagacggtagtcggatggttagatcgctcaagtaatttttg  
ttttgttgggtctcaggttttcagctgcccatttgatttcagtttgggcttttctta  
tctctaaaggcccaatttcatttaggtttagtttatttgatcattatccttactata  
aaggcttcgcctttcgagaaatttaggggtttcttctgtctgtctcgtcactcaggtt  
tgtgcctcaacgactgcttcaactctagcttgattcttcttcttctgtttatatgtat  
actgtacattagattattcttgtttctcgagcttctgctatagattttgattctttt  
ttttgttgtctttgtttcgtttccaggatcagatcttagctaaattgagacaagctc  
aaaatgaggtacttgacgcatctctacattcactgtttaattagagaacaatacgt  
ctctgaatcggtgattcagagacgtattgttcttctgtcatatgcaataagtttaatt  
agagaacaatacgtctctgaatcggtgattgttttttggatgtgcgttattgatagct

61

ttatgatgttaatagttctaggattgacacgaagttgttctgcagttttgcataaatg  
ctctttactaaggcctctaaatttggatgacaaatctaaatcttgcctcataaaaaat  
ttaggtgtattaagataagattattttgtatggtagtgtctataatgtgggttgttc  
atgttgaggttgtcaatgttgtgtattttgtttgttttagttaatttgccttaactct  
gttctttgtgggttaatacagtaagcttcagagtgaggccgttcgtgaagccatcac  
tactatcacagggaaatccgaggcaaagaaacgtaactttgtcgagactattgagct  
ccagatcggctctgaagaactatgaccctcaaaaggacaagcgtttcagtggtatctgt  
caagttaccacatatccccgtcctaaaatgaagatctgcatgctcggagatgcccc  
gcatgttgaagaggtgatataatcttttcatggaaattgatcattttgtgctctgttt  
cttgatataatggttttgtgctcatttcatttgggtggctctattagtttcatttgatg  
ttgtatatgtcttctgaatgtagatgcatgatgttttcggaatttgggtcattgttta  
tttaggcttcatttccttgcataattaaatatttgcttatttcatttctgtatcttttc  
gtaggctgagaagatgggggttggaaaacatggatgttgagctctcaaaaaagcttaa  
caagaacaagaaactcgtcaagaagcttgcaaagaaataccatgcttttcttggcctc  
tgagctctgtcatttaagcagattcctcgtcttcttggctcctggtcttaacaaggcagg  
caagttctggctacagctaatattccattgttcttctttacatccgttttgatttg  
gataggttttagtagtctatttcttttgtcaatgtctttttgatacaatgccaatcc  
tttatcctgtgagattatgcttctttgatgattcttaagtaacattcctttgcttta  
ctttacacaggaaaattcccaactcttgtagccaccaggaatccttggagtc aaag  
gtgaatgaaacaaaggcaacagtgaggtccagctgaagaagggtctgtgcatggga  
gttgcagttggtaacctttccggg

**Diaminopimelate Decarboxylase from Arabidopsis  
(clone#4159709)**

ccccgggtggcaaactgagatataagaggggaaggtgattttcatgcaaatttttttt  
tatttttttttgaatgaatgcaaaatttattcaaaaaaaaaaacctgggctacatc  
aagtacttcatttctgagtttttgaaaaaatctaaagacaacaaaagactttacaatt  
taataaaaaataataaaaaatactttatcactctcaacgaaattgttgatttaataa  
cgtatctcttggtaaaacagcgttttattttgacgaaattgttataaatgaataaaat  
gataatagaaactagtgtggtacgttaaaataacctctcatttggcaaaataacggtta  
tgtatcatgagttatgcatacgcagcgtgcttaaatagtgtgctttcaggagaaaa  
tatataccaagttatttgcgtgaaattaccacgcaaatctgaggttcgaatggcaaaa  
taaaaaaccaatgtcatttccttaattgtattaagggtcatttaataaaaattgtacac  
ttttttcacctgtaagcgttccaaagtgtagaatggataaactagaagggtcaaagg  
ataatattaataagcgaactcactttttgcccagtgatttcacttcttacatttgc  
ttgatatagtttacccaaaagtgtatatatttcccttatacaattgttctattttct  
ggattataagggggaataagaaaaaagaaaagagagagtataataataacttttata  
aagtgtgttagattctaatttgttaacgaaaagttcaaagtgaagaaaaaacgaaa  
aagtttttctgttttgttttatatctatagccaagaaagtttctcagatttacaaga  
agttaactgagaaaaacaaaaaaaaaacttatgaagcatgaaagactaattaacgag  
gtgattaattttgagacaaattaaacatcgaattaaaagtaaacatttggagggttta  
tatgttatatatgtgacatgataagtcgattcatgactaatgtatatctggaatct  
aacatgggaagaatagagaacgaagcagagccaagggtcaacttgcagacacgaatca  
acagattgtgaatgagaccaaatacaatgggtcataaacccggttgggtttaaaccggca  
agtcattccttgggtcaattccattcgttattccttcatgcaagaccctctgatacaa  
ccaaagactcccattacaatatcttttcgatcacgagctacttattttcaaatgtgt  
tacctctttcgtgactcttgtgttgtgtggttaagcctagtgcagatgtgtcggtat  
atataggcatacatatacaaatgcgacaaaataagtatattatattgtttaatttct  
atattccatttctatatgcatggctgggatttttgaccaaaccctaattcaagaat  
agaatccaaaagatgggatcaaagaatataatctaattgggctgaccacattttccga  
tttaattcgcatagttaatatctttccactactttatgcccgcagaaatttgaatt  
aagtaagacaaagaaatacagatataagatgggtcgtagaaaccagtagaggaatttc  
atttttcgtggataagtggaatattaataagagaatgggtctttactctttacagtgg  
gaaatgggaatagttagccattataatttcatcagattctatatatgcatgtttgta  
taagctaaaaataaatacgtttaagcattctcaaaaaaatttacaagttctagagac  
tctcttaacgtcggcaatttatattctactttacatgacactttcaggaaaagaaaa  
ctactactagcagatcattaaatttcttttcttttttgaatgaaccttagt  
tgtgggtttttatttttgttagctagaaacttcagtgtttttttccgccaatggtag

tgctttgatgatggtccggcccg

**Peroxidase from Arabidopsis (clone#4006885)**

ccccgggcaatcaaggtaacgaaggaggatcagcgaaaggatgggctatatttggagt  
tttttcctgctgtaagtaaatgctttgtgatcttccatgcggacatataactgaaga  
ataaactcaactcattgtgttctgggtgtgttcttctgatcagattcctcggtgcat  
ctgcacttttctgctgtgggggctttatttataaaacaagagtagagcgtgtggtaa  
tcctcatatcttctacaattccacttccattctctaattattctctcacgtgatat  
acacacactcaatcactgatgtactcgtatggatgcagcgtggaactgatgcattgc  
cggggatgtcacttctatcgggcttactagaaactgtaagtattacaagaaaactca  
aaaggattccatttatgcaaaatctaagagaaagctcactgtggtctttggttaca  
tttatggatctctcaagagacaaatgctatgtaagctaattgattttggtcttgata  
aacagggtgagtggaagtggacaaagctactcaagaactgaagacatcaacaatgctt  
ttgccaatgaagtctcatgggaccgctcttccgcatcttctactcaagcgacaacaa  
cacagagaccaagtgaagaacatattggtgcgatctaattttgtcaagtgcctcaca  
agaggtactgtttcaagccatgggtatggcacgcttgatctgcgatttctggtatt  
tgctttgtatgtttattttctacttctagaaagaggtaaaaaagttaatagcttca  
ccgtgagaatgttgttttaccagattcatgtgctatgatagaaaaagacaaagcaa  
acaagagttctttctttgcttaggttacaagaacaagagtatcgttataaagtcaac  
aaagattgaacatatttttgtcaaggagtggttagaatctcttccactctcttg  
cctttctcactaagacaaaaaaagacttggaactttgtctaaggttttgggtatatt  
attaaccaagtcttttgcaaaaagtaataattgtttttcgcatctctcttttagaa  
tttagtttaactataggtctttatattggttattactttctgaaaaatgatctgttta  
ttctattcacttgggtacctcgctttttatcttacttctacaaaaggattatcag  
tgaaagttagtctcttactctcaccttccgaaaaataaaacaaaaatatcgatacttc  
tagatcaaaccaagttgattaaaacatccctattccctacgattctgatcttgagat  
atattatcatgttaagatctaaattgacaagaaaactgatttttcatcttagtagg  
aaaaataattactattagtgatcatgattgtcgaccgtaagaggtgggttagttact  
ctccatctttcttgaagaagtcaagaagtcagaaattatatcaaattaaacatcaa  
tattagtaacatataatctgtatgggttttattgtttagaaaattccaatatttatata  
tcctagggaaaaagaagcttattcttcaaaattattgttatgagtcgttaaaatatgg  
ataaaaaatataaagtcataatattaaaaactcagtttgctttgcttttacctctcca  
agtctccaaagtc aaattatttttagttaattaaacaaaaaaggtttattagtc aa  
acttagcatgcaatgctgggtaccaaaccgaagcattagtccttttaattctcttt  
ttctccaataagtttttacaatttttaattgtttgcatttcccttgattatttatct  
tcatcccaatttagctaataaccaactccgtttcttattcttccaagtcctttcctat  
aaatacgttcttctccctcttattttcatatcactcaccacaaagtccttctcattt  
cctcatccccggg

**Mitochondrial Uncoupler from Arabidopsis  
(clone#4220510)**

ccccgggatgtgtgagtggaaggagaagaagagggaaacaaaggtatttattttagtc  
gagttttgttttgtgacgcggttttgtctgtgttcaatgttgacgaaacgagtgaga  
gagtgctgattattaaagaaaaccctaattaagtcagacccgcggttataaaaat  
agtcaaaaagtaggaaaacgcgtgtgtgagtgagacagagacagccattgtttgct  
ttatgggcttataagcgagacgtgttaattgggctttttcctttatggccgaaaaca  
aaagaaaagtcgcctgagagattcgaactctcgccgggcagagcccatgtacttagca  
ggcacacgccttaaccactcggccaaagcgacttggttgctatgagttagacaaaatc  
attaaaattctctattatgatttctcatagtggtgtgtatattgtggatctactaa  
aaattctttgttattattactttattttgtgaattagtttgataggttaagtacaa  
agtttaactttattatttactcaaaatttatcagatttaactgattttatattgttcc  
tttgggtatagacgtactatagtttttagaaaaaccataagattccttttatatttc  
atagagtgaagagatgagatgagatcttggctggagaagaaataagtttccacgagg  
aggactcttttttttgggtgaagacgaggaggaggactcttggttgatccagtcctt  
acgttagacatcgacccctacattttatttgctttctctatcaacatggcaggtaaa  
aatcttcattcaaccgaaccaacaaagtccttcccaataatattcaagcaccatc  
ctttgggaaactcatacatactacagtcctacactctttcattttctttcaacgctca  
acttaacaaatgatatagtctagttgtcaattatatgttttaattagtggtttcaca

63

tcaaattctggtttgatatttgatgacttttccggaacatctcaatgtcccgcaa  
atacaatcgtctatcatatataatcccgtacgttgtattcttatagatagaataata  
tggcgtgatctttataatataacatatagaatcgtgtagatttattttattttattt  
ttatatatcgcataaattgcaaaatacttatatatgtttgttatatatgatacccat  
tttatagttacttaaaaaaaggttaagcgataatatatatatcaactttttataac  
aaaaaagtataacacatggttaagaaaaataaaaaatgaagacatggtgtgacacgaa  
aatggcactaaatatacatatataatagatagctacaatatcccatcatacacactt  
ttttaattgactaataacataacttacacacttttttaattgactaattcataacttt  
ttatcattgtcaacatgcaaattcatatttccgttgaactattattcttattttgtt  
tttaaaagaagggcttcttggttaataaaaaatagatttccaaatgacgttagagcaa  
aaaaaaaaaaggttgtctggtctggttaaaatgaaaaagcaaagcgtcttggtatag  
aaaagtaataactgcctcctaatttcttcgtccttctaccgaagaatctctccact  
cttgccctctttcgaaaccctaaaccagaagcaccagattttttcaactttttccca  
gagaacaatagaaaacccaacttgtgctctctagggttttctttattccttctcatc  
tttgattttcttggtcatcttttgggaagcttaccaccagcgaaaaaattataa  
cttccatcgattcctggcttctctctctcgctctctctgcatgtgctaaatcgccgg  
actgatcctcactgtcacctctgttcccggg

**Stress protein from Arabidopsis (clone#6598614)**

ccccgggattaggggtttgagttgtcactggaaagaggtttgattgtgagtgatgat  
ggagagattatgaaggagtttgtgtgtatttatagaggagttagggttttgaggttt  
gatgagaagtaggtttgaagaagtttgttgttgcaacttatttagagttacttgtt  
ccacaaccacaagtaagattggtcacttctaagttctaactagaaacaaccatgaca  
catggagatttcagctaaccctagtttaattgtatatgtattatattttatttaaatat  
tataaaataaaaataaattttcacaaataaaaagaactacaaaaaagtgagaaaaataa  
tttgataaaacaaatttagaaaatttagtatatcaataaataaatttataatccgatgg  
ttttgccttttgggttggccttgggttgaacttcgatgagtgactatgtatagcgaa  
aacaattcgggttgggttgaatttttaaaaaatacaagcgacaatatctgatg  
agaataaggtgaaaagcaaataatatcagtttaattggaaatatttacttttttaca  
ctaataattttgttgggtcaaccaacaaatagatttaattaattatgggttatgagct  
tttatttgttgcgacagtatatatatgtttaaatagtgatattgcatggcggaaggt  
ccggaagcaacacatatctcctttttaatttttttttaacaagaataacatgttaa  
tttttttttgaatttaataaagaatacatatttctaatttttgcgtcagatagatga  
ttaagagtggtgtgttttttttaacaaacaaggaatacattatacatatttcatatt  
tctctcgacattgttgggttttttttaaaaaatagattaaagagtctacgaagctaagt  
agctaacgaagacttgaaatgagaagaagacgagaatcttttaataattttttgttaa  
gcgataatattttgaatttaataaatatagatttaaggaaataacaataacgcagat  
atcggttaagtcatagaaaaaagaacaacacaaacttacataaacatgtttcctaa  
tttgtaattggagttaaattccttcttttttttttttttttgatttggattccaatta  
gtaaagaactcaatgactataaataacctttaaccctctcattatttcttactatca  
attgattaagctctcgttcctaagaaagcaatagacgaacaagaacccatcgaagaa  
cacaaatctctcttgaagttgtcgataatgttagtacaccgttacttcgtccaaga  
cttttttgcggttccgtttcttacaacaaaggatttgggtaccattacttttgcg  
taactcctttttacatgtacgtcaaaaagtggttctcgtcgggttgaagaaacg  
accttcttaccacaaaaagcttatttttaaccgtctaaaaccggaaaaatctcaatc  
taaaccggatagcgttcatgagaaaccgattcaaacaccgagtgaagaagtagaatt  
ttttgatgggtccgtcaccaatgtgtgctgctccttcgccaagacatgtaccgattcc  
gatattttgtggtgtaaagatgatcaaagagctctcaaagctaagcacgacttgaat  
gagaagaagaagaccaattactcaattagattttgttttgggagcaattattgtct  
atttatctttgttttttagcaataaatctgtatccactaatcttcacagtacttgact  
aacaagaagtaaaagagttttcttattttccaattgttttttaactctgatactttttc  
ataattttacaatgtttgatgaaaaaaacattcaaacctaaattttcttttttggg  
tatgaattcaaacctgaattacttttgacgaggacccgacggtataaatagggtgat  
ctcccaacaaacaaaaaggttcccggg

**Pectinacetylsterase from Arabidopsis**

(clone#6671954)

ccccgggtggtggggacaatggatccggtctgcgtagcaacaaggctgaaaaagatta




FIG. 1




FIG. 2




FIG. 3




FIG. 4



Selectable Markers

pNOS / NPT-II / tNOS

pSU / Bar / tNOS

pSU/ Intron / Bar / tNOS

pUBQ3 / Intron / PMI / tNOS





FIG. 5

# Scorable Markers



<sup>1</sup> Construct useful for promoter analysis.

<sup>2</sup> Construct useful for high constitutive expression of genes of interest.

FIG. 6




FIG. 7




FIG. 8

AKK110P1  
SEQUENCE LISTING

<110> Mushegian, Arcady R.  
Taylor, Christopher G.  
Feitelson, Gerald S.  
Eroshkin, Alexey M.

<120> Materials and Methods for RNAi Control of Nematodes

<130> AKK-110P

<140>

<141>

<160> 139

<170> PatentIn Ver. 2.1

<210> 1

<211> 165

<212> DNA

<213> *Globodera rostochiensis*

<400> 1

```
gtttgagatt attgactttg catatttcca accaagttca tttgaccaat attttcctgc 60
taaacatagc aaaaatggtg aaaccgaagg tcggcattaa tggctttgga cgcattgggc 120
gcttggcggt gcgcgctgcg gttgagaagg acaccgttca ggtgg 165
```

<210> 2

<211> 342

<212> DNA

<213> *Globodera rostochiensis*

<400> 2

```
cgactacatg gtatacatgt tcaactacga ctcgacccat ggccgcttca atggcaaaat 60
ttcgacaagc gccggcaatt tggtcgttga gaaagagggg aaggccacgc acaccatcaa 120
ggtgttcaac ctcaaggacc cggccgagat caaatgggct gaggtgggcg cggaatatgt 180
gatcgagtcc accggggtgt tcaactacat tgagaaggct tcggcacact tgaagggggg 240
cgccaagaag gtgtcatct ctgtccgct cgctgatgca ccgatgtacg tgatgggcgt 300
caacgaggac aaatatgacc cggccaagga caacgtgatt ag 342
```

<210> 3

<211> 205

<212> DNA

<213> *Globodera rostochiensis*

<400> 3

```
gaagccggcc tcattggacg ccatcaaggc ggcggtgaag aaggctgccg aagggaattt 60
gaagggcatt ttgggttaca cagaggacca ggtggtgtcc acggactttc ttggagacag 120
tcgctcgtcg atcttcgacg ctggggcggt catctcgttg aaccgcact ttgtcaagtt 180
ggtcagctgg tacgacaatg aattt 205
```

<210> 4

<211> 167

<212> DNA

<213> *Globodera rostochiensis*

<400> 4

```
ttaaacgatt tattcacacg cacggagaaa tgaggattac ctaatttgat tgagtctttc 60
tcgtccattt gtcaattgtg gccctaaaga gggccgtttg ggtagtttt ttggtgttcc 120
ttctccttgc tggctcaacc accgaagccg tacagcgtcc ggccttg 167
```

<210> 5

## AKK110P1

&lt;211&gt; 41

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 5

catggccgtc acggtcttgc gcttcgcgtg ttcgcagtat g

41

&lt;210&gt; 6

&lt;211&gt; 79

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 6

gtttcccagg aaaactttca gcacggaacg agtctcctcg taaatgaggc cagagatgcy 60  
cttaacgcct ccacgacgg 79

&lt;210&gt; 7

&lt;211&gt; 168

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 7

cggcttgggtg atgccctgga tggtatcccg caagactttt cggtagcggtt ttgcgcctcc 60  
ctttccgagt ctttttccgc cttttccgcg tccggacatt ttgttgtaa atcagaagag 120  
cacagagagt aggagaaata ggaaattttg cctcgtgccg aacgtgcc 168

&lt;210&gt; 8

&lt;211&gt; 330

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 8

gacagtctcc gttctgggtta tgtgtcacac gcgcgaactt gctttccaaa tttctaagga 60  
atacgagcga ttcaccaagt acatgccggg agtgaagggt tccgtattct tcggagggat 120  
gccgataaag aaagacgaag aggtattggc taagaacacg ccgcacattg tcgtcggaac 180  
gccgggacgt cttttggcct taggacgcac tggacatctg aagctgaaag gcgtcaaatc 240  
ctttgtgctg gacgaatgcg acaaaatgat tggagatgcc gacatgcgcc acgacgtgca 300  
ggaaatcttc aaaatgacgc ctcaggagaa 330

&lt;210&gt; 9

&lt;211&gt; 136

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 9

actttgccgc gggagctgcg cgtcttctgc aaaaagttca tgcaggaacc aatggaggta 60  
tacgtcgacg acgaggctaa gcttacgctt cacggctctc aacaatacta cgtagactg 120  
aaggaaaatg agaaga 136

&lt;210&gt; 10

&lt;211&gt; 141

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 10

tattaaaata aaatacaaac aataatataa tggctgtttt ttctgtcatg tttcaagttt 60  
ttgttgttca tcactttctt cagcagcgac aatacggcca atccggtgaa agggccaaaag 120  
tcaatagctc gctcgggtacc t 141

&lt;210&gt; 11

&lt;211&gt; 141

&lt;212&gt; DNA

AKK110P1

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 11

aaccaggcac tctgttcac ttcggcatcg ctttttggca atgtcaacaa cactttgctg 60  
gccattttgt ttctacagca cacgcacacc gtcgtcttta cagcgttcac ctgcacaaa 120  
aagtagccgt atttgcgaaa t 141

&lt;210&gt; 12

&lt;211&gt; 37

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 12

gcgttggtg caagctgtac acaaggtcgc ccggttt 37

&lt;210&gt; 13

&lt;211&gt; 161

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 13

gcgcgttcca tcgcccgcac cacaaaaagt cccatcgctt catatcgtag cgcaaattgt 60  
ctttggtgca aatggcaaaa cggccaaaat aatggtcgaa gccgtacaca accgccaccg 120  
ccacagcgcc aacccacac caaatgcgaa atttatcgaa a 161

&lt;210&gt; 14

&lt;211&gt; 306

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 14

gaattcggtt gaggtataaa taaataataa atggcagcca acgaatcgct aaatgtggac 60  
agtttgatca ctcgattgtt agaagttcgg gggtgtagac cgggaaaaac agtgcaaatg 120  
gacgaatctg agatacgac tttgtgcatc aaaacacgtg aaattttgct gtcgcagcca 180  
atcttggttg agctcgaggc accttataaa atttgtggtg acattcacgg acaatataat 240  
gatcttctga gattgttcga atatggtggg tttccaccgg aagcgaacta tctatttctt 300  
ggggac 306

&lt;210&gt; 15

&lt;211&gt; 261

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 15

gcaaagcctt gagacgattt gtttgcgtct tgcttacaag attaaatatt ctgaaaattt 60  
ttttcttctt cgtggcaatc acgaatgtgc ttcaatcaat cggatttacg gattttatga 120  
tgaatgcaaa cggaggttcc tcaatcaagt tgtggaagac ctctactgac tgcttcaact 180  
gtctgccaat tgccgcttta atcgacgaaa agatcttttg ctgccacgga ggctgtctcc 240  
tgatttgcta aacatggcag c 261

&lt;210&gt; 16

&lt;211&gt; 151

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 16

gaattctttg agtgcattca gcgtttaatt ttttcgtatt ataataagca tggctcgcgg 60  
acccaaaaag catttgaagc gacttgacgc acccaaaaaa tggatgttgg acaaattggg 120  
tggcgttttt gcgccacgtc catttgccgg a 151

&lt;210&gt; 17

&lt;211&gt; 306

## AKK110P1

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 17

```
tcaagtacgc gcagtcgtac aacgaagcgc gcatgatctg caaacagcgg ctgatcaagg 60
tggacggcaa agtgcgcacc gagatgcgct tcccgtgcgg aataatggat gtgatctcga 120
ttgagaagac aaacgaaacg tticgtctgg tgtacgatgt gaagggccgt tttgtcatcc 180
atcgaattca aaagctggag ggccagtaca agctgtgcaa agtgaagaag caggccgctc 240
gggacaagca ggtcccctac attgtcacac atgacgcgcg caccattcgc taccggaccg 300
ctcatc                                     306
```

&lt;210&gt; 18

&lt;211&gt; 528

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 18

```
gaattcgcac aacgaattga agacttatgc ggcagaaaaa ggacttttgc caaagtgtga 60
ggagcaagca gacgaccttt cggattggct ttgttcgtcc attgggttgg agcatcgccc 120
gttcttaccg tatacaaacg ctgtaataaa tgaaacaatt cgattagtca atttgatccc 180
gttcaatctt agccatttgg cgcttgaaga tatgcaaatt ggcaatttta ttgtgaagcg 240
tgggacacca attgtaccgc aggtcagcag tgttctgttc gacgaaaaac tgtatccgga 300
gcccgatcgg tttttgcccg aacgctttct ggacgatgag ggccgtttga agaaaagcga 360
cgaaacttatt gcatttgggg ttgggaaaag gcaatgtgcc ggcgaaagctt tggcccgaat 420
gacacttttt ctgtttgccg ctaatttctt tctcgcctac aaagtctctc cgtccgatcc 480
actgaatcct ccaagcttga aaaagttggc ggattatctg ttacaca 528
```

&lt;210&gt; 19

&lt;211&gt; 335

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 19

```
gaattctttg agaaagcggg aattcgtttt tggctataaa atgattctgt gggccacgat 60
tttgttgatg gctttggaca ttgcgttcgg tggcaccaat caaatggaat ttgatcagtc 120
ggcgccgatg ttccccgact cccagttcat cgatttgatt tcgcgcgaca tcgaatcctt 180
ctccggccca ttgggcggtg gccataaatt tatgagcggc ggtgcccgtg agggcgctca 240
acagctaggc cccgaggggc cctttgagca gcggcaacag gtgaagagtg acaatgttct 300
ccccgcgtat tgcgagcctc caaatccctg tccga 335
```

&lt;210&gt; 20

&lt;211&gt; 52

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 20

```
ggacggctgc acggaacagt tcgagaacac tgccgagttt tcgcgcagct ac 52
```

&lt;210&gt; 21

&lt;211&gt; 190

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 21

```
gcttgtgtga ccaggagcac atgtttaact gtccgtcgaa gaacaaccgc gaggagtacg 60
agcaggatct ggagcaattg ctggccaaca acggactgca caaatcaatg attgccaaga 120
aattccatct ccgcggggcg gaggagccgc gccgtcgaaa acgctcttgt cgcccggctt 180
cggccaaccg                                     190
```

&lt;210&gt; 22

&lt;211&gt; 52

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis



## AKK110P1

<400> 22  
ccgctacaac ccctacctgg agggcgcccc gctgaagtca gtggccaaaa ag 52

<210> 23  
<211> 54  
<212> DNA  
<213> Globodera rostochiensis

<400> 23  
gaattccgac tctcaagggtg gaccacgccc caaccaacag caattgtcag ctgc 54

<210> 24  
<211> 77  
<212> DNA  
<213> Globodera rostochiensis

<400> 24  
ccgcacatgt cgaggcctcc atctttttggc actgggtcatc accttccgcc tactgctaac 60  
aacagaccgg aacagca 77

<210> 25  
<211> 439  
<212> DNA  
<213> Globodera rostochiensis

<400> 25  
gtcaatcaaa aacgccgact tcgattcctc agctgatggt cagtaatgcg ctaaaggggca 60  
tccattccgt ctcttctaca tcagcaacac aatcacattc cagccccagt tttatgacac 120  
acaacgtgca gcagcaacat gttgttgggt aacaacagca gcaacaacag aattttccaac 180  
aaccgcccgc cctatcgtag actcacagcc accaacaaca aaaacaacca ccacaagcgt 240  
cacagtcgat gttgtcaatg aaaagtggca atgttgtcgt tgttgttccg caacaatcgc 300  
agcagcacca ctaccaacag cggacactga cgccactgaa gcacacatcc gcatcctcca 360  
cgtccgatcg cttcgtcatc accaaaacca acaggggtgct tccactcccc tcgcagcaag 420  
gcgccacggc cactgatga 439

<210> 26  
<211> 539  
<212> DNA  
<213> Globodera rostochiensis

<400> 26  
gaattcgttt gagacacatc caattaatta atagttattg ttggcaatgg gacgagttcg 60  
aacgaaaact gtgaagaagg cgtcgcgcgt cattattgag aagtattaca ccaaattggg 120  
cctcgacttt cacaccaaca agcgcaattg cgaggagggt gccattatcc caagcaaacg 180  
gatgcggaac cgaattgcgg gatttatcac acatctgatg aagcgcattg agctggggccc 240  
tgtccgtggc atttccatca aattgcagga ggaggagcgc gagcgtcgcg acaattacat 300  
gcccgaatc tcttacctgg atgcgcagaa tcaccagatg atcagcaccg accaagagac 360  
gaaggatatg gcggaatttc tggggctagg cctcaacttg gaagtgaag ggccctttgac 420  
gagtggcggc gctggcgag gacgtcgttg agtcaggaca attggcatta ttgttgaaaa 480  
atcatcgatg tttgttcgc atttggatga taatgcgctg ataaattttt gttgatttt 539

<210> 27  
<211> 179  
<212> DNA  
<213> Globodera rostochiensis

<400> 27  
gaattcnaca gtttctgtga gtaatggcat ntcacactgc cggcatccaa cagttgcttg 60  
cggccgaaaa cgtgcggca gaaaagatta atgatgccc gaagcgaaaa gcacagcgac 120  
ttaagcaggc caaacaagaa gccacggcgg agatcgagca gtatcgnacg gagagggag 179

## AKK110P1

<210> 28  
 <211> 133  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 28  
 gcaaaattat ttgggcacgc gcgacgacat cgagcagcaa ataaagcgcg agacagaaga 60  
 gtcgctggag gcaatgaatc gcaatgtcgc ggcgaacaaa cagcaggica ttgtacgtct 120  
 gctgcagttg gtg 133

<210> 29  
 <211> 482  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 29  
 gaattcgtga aatcaaaagc ttttttaatt tatttacaca aaaaatgggt ccaccaccaa 60  
 ttcgcgtgtt ggtcactggt gccgctggac aaattggcta ttcactggtt ctgcaaatcg 120  
 caaaaggcga tgtgtttggc aaagatcagc caattgttct cgttctcctc gacattccac 180  
 cgatggccga agtactctct ggtgtccatt ttgaattgat ggactgtgcg ttggcaaac 240  
 ttgccggtgt ggaggctgtg accacggaag agcaggcctt caaggacatt gactacgctt 300  
 ttcttgctcg agcgtatgcc cgaagagagg gaatggaacg aaaggacctt ttggcggcaa 360  
 atgtcaaaat tttcaagtcc caaggcgaag cattggcccg cttttccaag cccgtncgtc 420  
 aaagtctctg tgggtggcaa cccggccaac acgaacgcgt acatttgcg ccaaatatgcc 480  
 gg 482

<210> 30  
 <211> 605  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 30  
 gaattcaaaag tgccgaaagc gttcaaaata attccggcaa tggcaaattg ggaacaaatt 60  
 ctgacacctta cttccccga aaaatggagt tcagcggcga tgtttcaagc aactcgtgtg 120  
 tttctgcca ccggcacacc gtcacaatgc caaagggtca acactttggt gctgttgcca 180  
 cgactccgtg atgagattga cgagtacaag aagctaaact ttcatttgta tcagtgcctg 240  
 tttaaagcaa tgttcaagcc ggccggattt ttttaaggga ttattttgcc tctttgcaa 300  
 tctggcactt gactctccg tgaagccatc atctttgggt ctgctctgcg aaagatttca 360  
 ataccgcaac tccacgcccgc tgcagcaatg ctgagcatag caaaaatgga ctactcgggc 420  
 gccatttctt ttatcctacg tgttcttggt gaaaaaaatt acacacttcc tttccgagca 480  
 ttagacggcc tcgtttttca ttttcttgga atgcgctcac atcagggcga gctgccagtg 540  
 atttggcacc agacactgtt ggcttttgct gagcggttac caaaagacat aagtgcagaa 600  
 cagag 605

<210> 31  
 <211> 112  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 31  
 ccattcccat catcaaatta ccccgattta ctgcggttt tgcgcggcgc cgagtcgagg 60  
 aatgaggaat gtgaagcaaa tgtgcccgtt tatgcgcgta atgatgaaat gg 112

<210> 32  
 <211> 105  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 32  
 gaattcgttt gagcatttat ttgacaaaat ctgaataaat ggccgtacca aaagaagtta 60  
 ttgacaaaat cgaggcgggt tacaagaagc ttcagggaagc gtctn 105

<210> 33

## AKK110P1

&lt;211&gt; 425

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 33

```

aagaagtacc tcaccaagga agtcgtcgat gcctgcaagg ataagcgac caagcttggg 60
gcgaccttgc tggatgtgat ccagtcgggc gttgccaact tggacagcgg agttgggggtg 120
tacgtccttg acgctgaggc ttacaccttg ttcaagccgt tggtcgaccc gatcatcaac 180
gactaccatg gtggcttttg tccgggcagc aagcagccgg caactgacct tggtagcggc 240
aaaacgcana tgctgaccgg atctcgaccc cgaggggaaa atttatcaat ttcgacacgc 300
gttcgttgcg gccgtttcct ttaagggata cccggttcaa cccgtgcttg acnaaaggan 360
aactacnttt ggagatggga aacnaagggtc nagggccggt ttctaacatt ttnaagggcn 420
atcct 425

```

&lt;210&gt; 34

&lt;211&gt; 581

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 34

```

gaattcgttt gagcgaagag ttttgtggtt gacaccgggt tatggacttt tagcccgtag 60
tccttgacgg tccaaagccg cgttcagttc cgtgccgtgt tttttaaag aggcggagag 120
tttgacggtc attccaagca gccaatatac caccataaac aaatacccc cccaatcga 180
tccccccct ccaattctc cgcattatc gcattatcaa ttctaatac cacaaccact 240
gcatcattcc tttcccgacc atacgatgct aagtgaact ttgaaaattg gcttcacgg 300
agccggaaag atggcccaag cattggcaag aggacttatc aattcggggc gatacccggc 360
agagaatttg atggcgagtt gtccaaagac ggacgaggct ttactggagc aatgcaaaaa 420
attgggaatc ggaacgacgc acgacaacac ttgggtcgcg cgagagaacg acgtcatcgt 480
attggcggtc aagccgatgc acatcagcaa agtgacgtcg gaaatcgac ccaatttccg 540
gagggaaatc ttgcttattt cattgattag gaattacact t 581

```

&lt;210&gt; 35

&lt;211&gt; 102

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 35

```

gaattcgttt gagaatttta ctttatataa ttgacgttta atcagcagcc ataagcaatg 60
cccatcaaag catccggaga aacattaagg aagtttattg tc 102

```

&lt;210&gt; 36

&lt;211&gt; 34

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 36

```

tgcaaatgat gcaaacccca cgcttcacaa gatg 34

```

&lt;210&gt; 37

&lt;211&gt; 100

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 37

```

tcatgttggt gccaaatctc gcttctggta ctttacgagc atgctgcgtc gagttaagaa 60
aacacacgga gagatcggtt cgtgtcaaga ggttttcgag 100

```

&lt;210&gt; 38

&lt;211&gt; 176

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 38

## AKK110P1

tgaagaactt cggaatttgg ctccgttacg attctcgtac tggacaccac aatatgtacc 60  
 gcgagtatcg ctgatgttac cgaggccggt gccgtgacct aatgctatcg cgacatgggc 120  
 gctcgtcacc gcgctcaggc ggatcgaatt caaatcatca aagtgcacac ctcaag 176

<210> 39  
 <211> 155  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 39  
 gaattccaag tttagggtat tgtttgttat acgatttctt acaaattgaca gaacaaactg 60  
 agcgcgcggt ccaaaaacaa ccgatcggtt ttctgaacga caagttcaga acgcaaggga 120  
 ttgggaagaa ggcattccaac aaggaccgtt actgg 155

<210> 40  
 <211> 35  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 40  
 tcctcgcgag gctattgagg gcatatatat cgaca 35

<210> 41  
 <211> 70  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 41  
 tggaaatgtg tccatccgag gtcgcattct cactgggggtg gtgatcaaaa acaaaatgca 60  
 gcggacgatt 70

<210> 42  
 <211> 85  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 42  
 tcgtacaaa atatcgtcgc tatgagaaac gccacaaaaa catgtccgtc cactgttcgc 60  
 cgtgcttccg agatgtctct ctccg 85

<210> 43  
 <211> 193  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 43  
 agttcgggtc aatgtgctca aggtgatcaa agcatcgggc tcgaagaaag cgttcgacaa 60  
 attctgagtc ggccaagcca accgcgaacg gtcatttgtt atggttccta attgttgctg 120  
 tttttcaatt atttgtgtta aatgactgaa ttatgatca acggtatact agtattcttc 180  
 tgaaaaagct cga 193

<210> 44  
 <211> 219  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 44  
 gaattcattt agatttggtt tgaagctaga aatctttatt ttgggagtca acgacaatgg 60  
 gaagacgtcc ggcggttgt tatcgtata ttaagaacaa gccgtatccg aagtcgcgct 120  
 tttgtcgcgg tgtacccgac ccaaaaattc gcatttttga ttggggtaga aagcgcgcca 180  
 ccgttgacga attcccatgc tgcgtgcata tgatcga 219

## AKK110P1

<210> 45  
 <211> 489  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 45  
 tccgaggcgc ttgaggctgc gcgaatttgc gcgaacaaat atatggtgaa gaattgcgga 60  
 aaggacgggt ttcatatgcg cgtcagaatc catccatacc atgtaattcg catcaacaaa 120  
 atgttgctct gcgctgggtc ggaccgtctg cagactggga tgcgtgggtc gttcggaag 180  
 cctcagggac tcgtggcgcg tgtcagcatc ggtgatatgc tgatgtcagt gcgtattcgt 240  
 gaccaacacc aagctcacgc attggaggcg ttccgctcgg ctaaattcaa gttccctggt 300  
 cgtcaataca tcgtcttgtc ccgcaagtgg ggcttcacca aattcgatcg cgagggtatc 360  
 gagaaatacc gcaaggaggg ccgtgttatc cctgacggtg tgcatgcaa gttactcaag 420  
 caacacggac ccgctgaagg agtggtctaa gaacccatt taatcttctg tttgtcttgt 480  
 gactcttgg 489

<210> 46  
 <211> 101  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 46  
 gaattccccc gctcgagccg gggttgacgat gtcctcctcc acctcctctc actgcgttcc 60  
 gtcctccttc agccggaaat tgttctgtg gctgttgccg g 101

<210> 47  
 <211> 485  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 47  
 tccaccaaaag tccattcgct gtcgccagtc cattttattcc acaaaaagat gattccgctc 60  
 tcgttccgat gacgtcgttt ggccaaccgt tgcccccgct accgctttca ctggtgcca 120  
 acccgccgct ttattttgtg ttcccagaaa acttgccgtt ggagcggccc ttcgacgagc 180  
 aaaacgacgg ctccgaggag gaattagccg aagaagcgat gggaacgaag gcgaagagg 240  
 cgcaaacggt cgtccgattc ggcaaaaagg cgcaaacatt tgtgcggttc ggaaagcgtg 300  
 cacaacatt tgtagcctc ggaagggaca cgcaaggca attcgatggg aaaatgcaa 360  
 gtgaacagca acagaaaaag gcttaaagca aacggcgcg acttttctt taatgaatgc 420  
 gcgcccaccg catgacaatt cttttgtgta atgtgttgcg atttttatga tcggtaaatg 480  
 taaca 485

<210> 48  
 <211> 651  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 48  
 atctgttcaa gggactgttc ggcaagaagg aaatgcgcac tctgatgggt ggggtggacg 60  
 ctgctggaaa gacgaccatt ctgtacaagt taaagctcgg cgaaattgtc accaccatcc 120  
 caacaattgg cttcaacgtg gaaaccgtcg aatacagaaa catctcgttc actgtttggg 180  
 acgtgggtgg tcaagacaaa attcgtccac tttggaggca ctactccag aacacgcaag 240  
 gactgatctt cgtcgtggac agcaacgac gcgagcgtgt gggcgaggcg cgtgaagagt 300  
 tgatgcgaat gctggcggag gacgagttgc gcgacgcggt gttgctgggt ttcgctaaca 360  
 aacaggattt gccgaatgcg atgaacgccg ccgaactgac agacagactt ggactgcaca 420  
 acttgcgaaa ccgcaattgg tacatccagg ccactgcgc gacttcgggc gacggactct 480  
 acgagggact ggactggctg agcaaccagc tcaagaacag aggctaagct gggttggtgt 540  
 ctgttgact tgcccgcgga attgatgac attgaattta tttgtgtgt tgcgcgcgca 600  
 gctcttttgt gggacgtccg attaatattt ataattattt tattccgtgt t 651

<210> 49  
 <211> 660  
 <212> DNA  
 <213> Globodera rostochiensis

## AKK110P1

<400> 49  
 gaattcccaa gtttgagatc aattcagttt cacttagaca aaaatgccgc cgaaattcga 60  
 cccaaactgag atcaaaatcg tgtacctgcg ttgctgctgg ggtgaaattg gtgcaacatc 120  
 tgcacttgca ccaaaagtgg gcccaactgg attgtcgccc aaaaaaattg gtgaagacat 180  
 tgcgaaggcc acacaggact ggaaagggct taaggttacc tgcaagctga caattcagaa 240  
 tcgtgtcgcc aagatcgacg ttgtcccatc ggccgcctct ctgatcatca aagagttgcg 300  
 cgaacctccg cgagaccgca aaaaagtcaa aaacgtgaag cacaatggca acctgaccat 360  
 cgagcaagtg atcaacattg cgcgtcagat gcgccctcgt tcaatcgcac ggaagttgca 420  
 gggcaccgtg aaggaaattt tgggaaccgc ccagtcggtt ggctgcacca tcgatggaca 480  
 acatccgcac gacattgtgg acgcgatcag agggggagac atcgaaatac ccgaggaata 540  
 aagaaaggac ggcgcctccg atttttgtgg gacggacatt ggggaatttg ggtgaatgag 600  
 ttgccaatth cattcattca tcaattgttg ttattgntgg tacggataaa tttgtaattg 660

<210> 50  
 <211> 625  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 50  
 gtgccggaac agacgctcga ggaggttagc cgtctgcagc ggacgagctc cttgttggaac 60  
 gtggcaatcc gggacggcgt cccctacccc ccactgcctc ctacaaaccg atccccgaa 120  
 tacatgaaca tgctgaccgg ctcttctctc gtgccaaatt tccgcatcta ctggggcgcc 180  
 atcggaccgt acagaccttc gttgcccggt tacacttaca acacttacca cgggtacttc 240  
 ccctaccgca actaccgagg ctacaccttg gcgaatgctt actggtacga ccgatactat 300  
 tacttctcgc cgctgtacaa acgaagcatg ttccccaccc gcttcaaaca ttgtgactat 360  
 aaagcgaacc cgcactattg gcactacccg cacacctttt gggactatcc ctaccagggc 420  
 aaatggttcg actacgacaa ccctcccaat taccggccct actacaacca tcgccttaac 480  
 ggatattgctc ggccgtatca ctaccggtcc catgcgctgg ccacccggt caattaccgg 540  
 gaaggaatgg tcaggaaacg ggtctgacaa atcgaactgc tccaaattga cgtggtccgc 600  
 attcgaaaga agacgaaaa agctt 625

<210> 51  
 <211> 402  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 51  
 gaattccaag tttgagcaac attttgaana tgaccgaagc caaaaaactt cccgaggtgc 60  
 cggaactttt gctcaagcga cgcaaaatca gagctgcgca aaaggccgca aaagcaaga 120  
 acaaataggag ttctatcaaa aaagcacgga ccaagaagggt ggaaatcttc aaaagagccg 180  
 agcagtatth ggtggagtag cgtcagaagc aacgccaat gcttgcgctg aaacgtgaat 240  
 cgaagaaagt cggaattat tatgtgccag aagagcccaa actcgccttt gtggtccgaa 300  
 tcaaaaggcat caataagatt catccgcgctc ctgcgaagggt tctgcagctt ctccgcttgc 360  
 gtcagatcaa caacggcgtt ttcgtaaagt tgaacaaggc ga 402

<210> 52  
 <211> 433  
 <212> DNA  
 <213> Globodera rostochiensis

<400> 52  
 ccgaccgta catcgcttgg ggttatccga gtcagaagat catccgtcag ttggtctaca 60  
 aacgcggtta cgccaaagag aagggacagc gcattccaat aacggataac aacattgttg 120  
 agcgcagttt gggcaagcat gacgtgattt gtgtggagga tatgatccat cagatttgga 180  
 ccggtcggac cgcacttcaa acaggtgacc aacttcttat ggcctttcaa gctgagcaac 240  
 ccggtgggag ggttcaagaa gaagtccaat cactttgttg gagggaggcg attatggaaa 300  
 ccgagaggac caaatcaaca aattattgga aagaatggtc taatggaagg gaagcgana 360  
 aagaaaggaa attgnggcgt ttttctgttg ttgttttgac gataaattgt taactccaaa 420  
 aaaaaaaaaa aaa 433

<210> 53  
 <211> 768  
 <212> DNA

AKK110P1

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 53

```

gaattcgttt gaggtcaaac tttattagcg tatttaacaa tgtccgaagg aggagcgaaa 60
aagagtagca gcggtgccaa ggggggggtt gatgtcaaga aatttgcgat cgatcttgcg 120
tccggtggtg ctgcgggcgc tgtctccaaa actggtgttg ctccattga acgtgtcaaa 180
ctcttggtgc aggtgcaaga tgcctccgct cacatcactg ccgacaaacg ctacaaaggc 240
attattgacg tgcttggtccg tgtgccgaaa gagcagggct ttctgtcact gtggcgtggg 300
aacttggtcca acgttatccg ttatttcccg actcaagcgc tgaacttcgc cttcaaagac 360
acctacaaac gcatctttac ggagggactg gacaaaaaca agcagttctg gtcgttcttc 420
gtcatgaatt tggcctctgg aggtgcgggc ggcgccacgt cgctgacctt tgtttatccg 480
ctgggacttt gcccgtacgc gtttggtccc tcgatgtccg aaaagctggg tcccgcgagt 540
tcaacggttt ggccactgc atcgcaaaaa tcttcaagtc ggacgggtccc atcgggtcttt 600
accgcggtt cttcgtctcc gtccagggca tcatcattta ccgcgcgcc tactttgagt 660
gctttgacac cgcaagatg attttcgcgc cggatggcaa gcagatgaat ttcttcctca 720
catgggcatc cgctcaggtc gtcaccgtgt cgtccggtgt cctctcct 768

```

&lt;210&gt; 54

&lt;211&gt; 338

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 54

```

gaattccagc agattaattg gaatggctga gaacatcgaa gagattcttg ccgaaatcga 60
cggctcccaa attgaggagt atcaacgctt tttcgacatg ttcgaccgcg gaaagaatgg 120
ttacattatg gccaccctaa ttggacaaat tatgaacgcg atggagcagg actttgacga 180
aaagaccctc cgaaaattga tccgcaagtt cgacgcggac ggttccggca aactggagtt 240
cgacgagttc tgcgcgttgg tgtacacggt ggccaacact gtggacaagg acactctgcg 300
aaaggagctg aaggaggcat tccgactctt tgacaagg 338

```

&lt;210&gt; 55

&lt;211&gt; 267

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 55

```

gaaattgcgc ccgatctcag cgacaaggat ttggaggcgg cggtcgacga aattgacgag 60
gacggcagcg ggaagatcga attcgaggag ttctgggagt tgatggcggg cgaaaccgac 120
tgagaaaaga gcaaatcgat ccaaatccaa acggaccgtt cccatttcac ctccatccgt 180
ccgtcgtatt attatatatt ccagtggaaat tttcccatta aaattcgggtg aaagtaaaat 240
aatttgacga aaaaaaaaaa aaaaaaa 267

```

&lt;210&gt; 56

&lt;211&gt; 597

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 56

```

gaattcgtg gacacttcgc atccggagta cagccacgag cagagcatcg accagaccag 60
catcccctac cagatgggtt cgaacaagta cgcctcgagc aagggcagta ccggctttgg 120
acagccccgt tgggagggtg ttgaccgctc catctcgtag cagaaccgca agtcgcaagg 180
aatggttcgt ctacagtcgg gtaccaaccg gttcgcctcc caggcgggca tgaccggctt 240
cggcacaccc aggaacacca cctatgaggc ggaggcaggc gagctgccct acgaggacat 300
gaagaagtgc gaggcgatca tccgtccca ggccggttgg aacaagggag actcgagaa 360
gttgatgacc aacttcggca cgccccgtaa caccaccacc aagggtcaaa tggaagaatt 420
ggcggaaaatt ccggaggaca ttttgctgaa aggacacggc gaggtgcgcc tgcagtcagg 480
taccaaccgg ttcgcgtccc agaagggctt cgtcgcgttc ggtaccggac gtgacgtgtg 540
ccgtgagggg gtgaacgtga acgtgctgcc gggcgacttg gagccgcttc cggagga 597

```

&lt;210&gt; 57

&lt;211&gt; 80

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

## AKK110P1

&lt;400&gt; 57

ggcattgtgc gtctgcaagc cggtacgaac aagttcgact cgcagaaggg catgaccctt 60  
 ttcggtacgg gcccgtcgtg 80

&lt;210&gt; 58

&lt;211&gt; 513

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 58

gaattcgcga caccgctcac atcgcgtgca aattcgcgga acttaaagag aaggtggacc 60  
 gncggtctgg caagaaagt gaggacaacc cgaagtcgct gaagactggc gacgccggaa 120  
 ttgtcgaact gattccgacc aagccgatgt gtgtggaggc attcactgac tacgaccgcg 180  
 tcggccgttt tgctgttcgc gacatgaggc anactgttgc cgtgggcgcg atcaaatcag 240  
 tggagaagac ggaaggcggg ggcaaagtga ccaagccagc gcagaaggtc ggcgcgactg 300  
 gtggcgggaa gaagacatga ccaaggggag gggcgggtcc ctaagggcca accgtcgacg 360  
 aaaatgacgac caacctcttg tttatcgttg tcttattcag ttccttcac ccgtctctat 420  
 ccatattgtc gttgcgttgg ataattgttt atttttgtt attgtcctgg ttggaaaata 480  
 aatttggtca attaaaaaa aactcgtgcc gaa 513

&lt;210&gt; 59

&lt;211&gt; 393

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 59

gaattcgttt gagcgaaaaa aacatactat acaatggcaa caactgagaa gcctcaggtg 60  
 gttcaacagc ccgtgcaggt ctttggccga aagaagacag caacagccgt tgcgtttgca 120  
 aaaaggggca agggcttgat caaggtcaat gggcgtcctt tggactacat gcagccggag 180  
 attctgcgca ttaagctcca ggagccaatt ctcatgttg ggaaggacaa atttgaggga 240  
 atcgacatac gaatccgcgt caagggcggg ggacacattg cgcaaattta tgcaattcgc 300  
 caagcactgg ccaaggcact ggtcgtttc taccagaaga atgtcgacga gcagagcaaa 360  
 aaggaaactga aggagcaatt tgttgcttac gac 393

&lt;210&gt; 60

&lt;211&gt; 154

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 60

cacgagccaa agaaattcgg tggacccggg agctcgcgct cgctaccaga atcgtaccgt 60  
 taagaaataa tttgtagat caaatgttt gatgatgat cttgtttttg ttgttgataa 120  
 aaaaaattta taaaaaaaaa ccgccgatac tgac 154

&lt;210&gt; 61

&lt;211&gt; 666

&lt;212&gt; DNA

&lt;213&gt; Globodera rostochiensis

&lt;400&gt; 61

gtattccaag tttgagcgat cagagttcct caatctatta tcaactgttt tccatcaacc 60  
 aactgtcatc atgcaaattt tcgtcaagac gctcaccggc aagaccatca ctctcgaggt 120  
 cgaggctagc gataccatcg agaactgtaa agccaagatc caggacaagg agggcattcc 180  
 gcctgatcag cagcgtctga tcttcgccgg aaaacagctt gaagacggac gcaccttggc 240  
 cgactacaac atccagaagg agtccactct ccattctcgt ctgcgtctcc gtggcggaat 300  
 gcaaattttc gtcaagacgc tcaccggcaa gaccatcact ttggaggctc aggcagcga 360  
 caccatcgag aacgtgaagg ccaagatcca ggacaaggag ggcattccgc ctgatcagca 420  
 gcgtctgatc ttcgccggaa aacagctcga agacgggcgc actctggccg actacaacat 480  
 ccagaaggag tccactctcc atctcgtctt gcgtcttcgt ggaggagaga actgaatcgc 540  
 gggctgatgg aaagatgacg aatatgatgt ctattcgatg acttgtctct ttcgatataa 600  
 ttgattgtgt tccatttgtc ggtcatcaaa tctttatgac cccctcattg ggcattggaac 660  
 gataaa 666



## AKK110P1

<210> 62  
 <211> 213  
 <212> DNA  
 <213> *Globodera rostochiensis*

<400> 62  
 gaattcgttt gagaaacttt ttcaaccatt cattcaaag tctcatcaag tgacacgggc 60  
 agcactcaac cacgggacgc gtgtactgag cgtgttgag aagggtcaag tggctctgctg 120  
 gtttgaggag acacattcgt tcgcgcaagt ggctcgaaga taccgggcag aatttggtat 180  
 ggaaccaccg cagttggacc aagtgaagaa gtt 213

<210> 63  
 <211> 488  
 <212> DNA  
 <213> *Globodera rostochiensis*

<400> 63  
 agcaccggct caatcctcaa tggcacaacg acggcattct ccggcatagg agacggagtc 60  
 ggtcttgagg aacaacagcc aattcccgtc gtaagcgatg cgggactgga tgcggaagaa 120  
 cagctgagaa tggccagaat gtgagccgga ggacctgaag atttatgaac gaaattttcc 180  
 agtgaagtgg accaacgctc ttcgacttta tctgctttgt gtaaagtgtg tagaatcggc 240  
 ttccaattca aaggcttttc attccccaac ttttattttt gcgcaaaaaa tttcttagga 300  
 taagcgtgaa taattttattg atttggtttt tctttctttt atctccgcct cgaagtcgca 360  
 agtgttcctt ttggcccgtt cccttttgtt ttgaatgtta ttccattccc atcccctcac 420  
 tttctcatat ttgtgacatt cagctgcatt gttcgactcc catttaaaag ttgagtgaag 480  
 tgcgattg 488

<210> 64  
 <211> 249  
 <212> DNA  
 <213> *Globodera rostochiensis*

<400> 64  
 wccrgakbng aacahcdkdg vhwatnvcbn gschvbwagc rngtcsvddb wgnhnsswtg 60  
 gkgdyrbwnt msnwrmanrg artsstsgaa tccccaagtt tgagagtaaa tattatttagc 120  
 taataaatggc agtcggaaag aataagagaa tgggcaaaaa gggagccaag aagaaggctg 180  
 tcgatccgtt cacacgcaaa gaatggtacg acatcaaagc gccggcgatg ttcaacacatc 240  
 gaaatssts 249

<210> 65  
 <211> 362  
 <212> DNA  
 <213> *Globodera rostochiensis*

<400> 65  
 wcbcrbhdby ytsgcrsnck tbdshbcysy gcdwkmtnvk hscngdckty nyykkkvbmr 60  
 ntmsnwrman rgartsstsg tcaaccgtac tcagggaacg cgcatttcga gcgactttct 120  
 aaaaggccgc gtttacgaag tgtcactggg tgaccttaac agcactgacg ccgactttcg 180  
 aaagtccgc ctgatctgtg aagaggtaca gggcaagatt tgcctgacca actttcacgg 240  
 aatgtcgttc actcgggaca aactgtgtc tattgtcaag aagtggcaca cgctcattga 300  
 ggcgaatgtg gcagtgaaga ctaccgacgg tttcatgctc cgactctttt gtatcggtss 360  
 ts 362

<210> 66  
 <211> 128  
 <212> DNA  
 <213> *Globodera rostochiensis*

<400> 66  
 aatcaaatta agaagacgag ctatgcaaaa gcctctcagg tgcggatgat tctgtgcaaaa 60  
 atggtggaga tcatgcagaa agaggtctct tccggcgatc ttgaangaaa gtagtcaaca 120  
 agcctgat 128

## AKK110P1

<210> 67  
 <211> 502  
 <212> DNA  
 <213> *Globodera rostochiensis*

<400> 67  
 gaattccatt aaaaaactaa acgaacaaat ctaaagatgg ccaccgaagt ggaggaaaat 60  
 gttcctacgg ttgacccatg ggggtgctgtg gaggaagtgg gtgggtgaaga gtcgatgcag 120  
 ttggtcagcc ttgacgttac cgaggtcaaa ctgttcggaa aatggtcctt taacgatgtg 180  
 gaagtgtccg acatttcgct tgtggattat attgcggtga aggaaaaggc ggccaaatat 240  
 ctgccgcaca ggcgcggccg ttaccaacag aagcgcttcc gcaaggccac ctgtccggtg 300  
 gtggaacggt tgtctttgtc aatgatgatg cacgggcgga acaacggaaa gaaactaatg 360  
 gcggtgcgca ttgtgaaaca ccccttcgag atcatcacct gctaccggag agaaccaggt 420  
 ccaagtgttg gtcaatgctg tgataaacag tgggcccnc gaagatttca cacgtatcgg 480  
 acgtgcgggc actgttcgtc ga 502

<210> 68  
 <211> 519  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 68  
 gcaaaactttt atcaataaaa aaatttatat ttgccaaaca aatttatgaa taaaaattca 60  
 ttaatcatta aaactacatt taaaatatac tttttagaga atgtcgtcta aaatattctt 120  
 ttctcccctt tatgcatcta tctaaccaga cttgggaagca atatggctaa tcaagtcaac 180  
 aatacggcag gaatacccaa actcgttatc ataccagcta accaatttaa caaaatgcgg 240  
 gttgagaacc ataagagcct cggcgctcgaa aatagacgaa tgagtgtcgc caagaaagtc 300  
 ggtagaaca acctggctct cagtatatcc aagaatccct ttaagctttc ctccgaagc 360  
 agtcttaatt gcattcttaa tagcctcctt cgttgctggc ttctccaaac gaggagtcaa 420  
 atcaacaacg aaaacgtttg ggcgtcggca cacgaaaagc catttccggt aagcttccca 480  
 tccaattcat ggattgacct ttccaacagc ctttgcagc 519

<210> 69  
 <211> 218  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 69  
 ttgattcttt attagtggac aatgacggaa gaccagaaga agttgccgat ggtgcctgag 60  
 actgttttga agcgaaggaa agttagggtc gctcagcgtg cttctctact caagaataaa 120  
 ttggagaata ttaagaaggc taaggttaaa acgcaagtta tctttaaacg tgctgagcaa 180  
 tacttgattg catatcgacg taagcaaaaag caagagtt 218

<210> 70  
 <211> 293  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 70  
 taagaaagca gggaattttt atgtcccaga tgaacctaaa cttgcttttg ttgtgcgtat 60  
 taagggaatc aacaagggtta atttaaattt gctataaagt ttaggatggg tttagacaat 120  
 tcttctcttt taatgcittc taactttttc aaaaaagtta tgattttatc acccataaat 180  
 ctacaaattc ttttaattat cagatccatc ctcgtcctcg aaaagttctt caacttttcc 240  
 gcttgcgta aatcaacaat ggagttttca ttaaattgaa taaagctaca atc 293

<210> 71  
 <211> 422  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 71  
 aatgcaatta agactgcttc ggaaggaaag cttaaaggga ttcttggata tactgaggac 60  
 caggttgttt ctaccgactt tcttggcgac actcattcgt ctattttcga cgccgaggcg 120  
 taagtittga ttttctaaga ttatatttaa ctttttaaat ttttcagtct tatgggtctc 180

## AKK110P1

```

aaccgcatt ttgttaaatt ggtagctgg tatgataacg agtttgggta ttcctgccgt 240
attgttgact tgattagcca tattgcttcc aagtctgggt agatagatgc ataaagggga 300
gaaaagaata ttttagacga cattctctaa aaagtatatt ttaaattgtag ttttaatgat 360
taatgaattt ttattcataa atttgtttgg caaatataaa ttttttattt gataaaagtt 420
tg                                     422

```

<210> 72  
 <211> 374  
 <212> DNA  
 <213> *Meloidogyne incognita*

```

<400> 72
atctgagcat aaggaaactt ggcctcaagc tatagagcag accgattatg tggcaccgac 60
tgagccagtt aaactggact tcaacgttcc gcttattagt gattgggctg ctgcttctga 120
gtggcctcaa gaagaggaag ctgaggttgc acctactgca ccaattggct agccacagcc 180
tcaacagcag caaactcaac aaggagggtga ttggaactct ggtactagtg gatggtgaag 240
ggcaggaaaa ttgatagaaa gagaaattat tatggaataa atgtaatcaa tgttgttgtc 300
tgatttattt gttacatata caacaagttt tattttgttg tttatttaat aaaagttgtt 360
aattaaaaaa aaaa                                     374

```

<210> 73  
 <211> 120  
 <212> DNA  
 <213> *Meloidogyne incognita*

```

<400> 73
tttttttttt tttttcttca tcaatatttt gaagtgaaga accagaagta gttgcattcg 60
agctttcaaa ttttgttttt tgattactct ttaaacaaga ttcaactgat ggatctactg 120

```

<210> 74  
 <211> 369  
 <212> DNA  
 <213> *Meloidogyne incognita*

```

<400> 74
gtctaacc aa tctagagcta ttcgggttcgt ctgtctgttg attattagat gttgattgaa 60
cagcactagt ctctgatgta gttttcttca atctcatttt taagtgatgt agaggaagtt 120
tagaattctg attgctatcg tcttctttct ctctctttta tggctttttc aattttatctt 180
cttccttttc ttgtccattc ttttcttcat tcttttcaaa aggctcagga aatttttaatt 240
cagaccgct ccttttaact gctgtatcta aagaaaaccc tctaggcaac gtcccagttc 300
cactcaaatt caattttgtt aaatttttgc cagatctaag tccttcttcc ttttgaacga 360
attgaactg                                     369

```

<210> 75  
 <211> 529  
 <212> DNA  
 <213> *Meloidogyne incognita*

```

<400> 75
ttttgttttt tttttttttt ttatcagaaa aaagtttaatt cagaaaaaaa aattaaaaca 60
aatctaaata aggctctatt ctaagtttat atttttcttt tacataaacc gtcaaccctc 120
caagtttttc aatgcttgga ggttttaatt gatcctctgg taataatttg taggctagaa 180
aaaagtttgc agcaaaaagg aaaagcatca ttcttgctaa ggcttctcca gcacattgcc 240
ttttccccc accaaaaagct attagctcgt cagctttttt taatttccct tcattgtcta 300
tataacgttc aggtcaaaa ttttggggat ttgggtatat ctttggatca aaaagaacat 360
ccgatacttg gggatcata aatgtacct taggcaacac aaactttcca acattcaaatt 420
cttccaaggc taaatgcccc aaattgaaag ggactaaatt aacgagtctt aatgtttcat 480
taacaacagc atttgtataa attaatttag gtctgtgttc caaactaat                                     529

```

<210> 76  
 <211> 449  
 <212> DNA  
 <213> *Meloidogyne incognita*

## AKK110P1

<400> 76  
 agtttttttt tttgaataaa agactttttt ttattaaaaat ggcttcgcaa actgcaggaa 60  
 ttcaacaatt acttgcagca gaaaagcgtg ctgcagaaaa gattaatgag gcacgtaaaa 120  
 gaaaggcaca acgactttaa caagcaaaac aggaagcgca agctgaaatt gacaaatata 180  
 gagaggaacg tgaaaaacgt tttaaagagt ttgaacataa ttacctcggc gctagagatg 240  
 atattgctgc acaaatataag cgtgaaactg atgagacgct taatgaaatg actcgtagt 300  
 ttgctgctaa taaacagcag gtaattgttc gtctacttca acttgtctgt gacattcgtc 360  
 cagaactgca tcacaattta caacttcaac ttaagcttaa tgaaaagcct gcctaatttg 420  
 tagttgattg attataaaaa tgaaattga 449

<210> 77  
 <211> 643  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 77  
 atttatattt gaacaaataa tttacaaaaa aagtatggct cgaggaccaaa agaagcattt 60  
 gaagcgtttg gccgctccaa agaattggat gttggacaaa ttgggtggag tttttgcccc 120  
 acgtccccatg tgcgggcctc acaagcttcg tgaatcgctt cctcttattt tgtttcttcg 180  
 taatcgctcta aaatatgcac aatcttataa tgaagctagg atgatttgca aacaacgtct 240  
 cattaaagtt gatggcaagg tgcgtacaga aatgcgcttt ccagctggat ttatggatgt 300  
 ggtttccatt gagaaaaactg gcgaagtctt tcgtcttctc tatgatgtca aaggacgttt 360  
 cattactcat cgcatacaaa aggaagaagg tcagcttaaa ttgtgcaagg tagtaaagca 420  
 agcgattggg ccaaaaacaa ttccttataat tgttactcat gatgcccgtat ctattcgtca 480  
 tccggatcca cacatcaagg ttgacgacac tgttgctgtt gatataaaca ctggaaaagg 540  
 tacagatcac attagatttg attctggtaa tgttgtatg attactggtg gtcacaacat 600  
 gggacgtgtt ggtattgggt gacatcgtga acgccaccct ggt 643

<210> 78  
 <211> 584  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 78  
 atttcctcta aaaatgaatt taaaagaaca acaaatatat ttaaattattc aattattatt 60  
 ttttattttg gctgtcagta gttttttgac aactaaggga agtgaagtaa aacaacgaga 120  
 aaataataaa ttggaatata ataaaaatga aattgagagg caaaaagagc aattaattcg 180  
 agatttgatt gccctcttaa cacgtgaaag gcaatattca cgagattggc aacaatcaca 240  
 acagcaacaa aatttcatta acagtitttg ccttcccca catttattcc cctcttcagg 300  
 cattgaatgg ccccaacaac aacaaaaaat atttttggaa gaagggggaag tagaagaacc 360  
 ttttagaggaa aatgagaagg aaaaaagagc acaaactttt gttcgtttcg gaaagagagc 420  
 acaaacattt gttcggtttg gaaaaagggg acagactttt gttcgatttg ggagagattc 480  
 aaaaacatcaa cataacttgt cagatcagaa gcagttaaaa actgacaaac aataaaaaatg 540  
 atgaattatt taaaaatttt tttaatgata ttttaattaa aatt 584

<210> 79  
 <211> 556  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 79  
 atcaagcatt aaatatgcag atttttgtaa agactctcac cggaaaaact attactctcg 60  
 aggttgaggc ttctgatacc attgagaatg ttaaggcaaa aattcaagat aaagagggtg 120  
 tcccgcctga tcaacagcgt ttgatctttg ctggtaagca acttgaagat ggacgaacct 180  
 tggctgatta taacatccaa aaggagctca cacttcactt agttttacgt cttcgtgggtg 240  
 gaaagggtca cggttcattg gctcgtgctg gaaagggctg tgctcaaact cctaaggctg 300  
 aaaagcagga acataagaaa aagaagcgcg gccgtgcttt ccgtcgcat caatataacc 360  
 gtcgcttcac caatgttgct acttctgggg cgggacgccg tcgtggccct aactccaacg 420  
 ctgcataaga gaatggctgt atcttgatga atgtatgggt atataatcaa tttaatatcat 480  
 tcgactntat gaagttttct gttattcaag ataaatcttt ttgttgaaaa aaaaaccaag 540  
 tttgagatca gttact 556

<210> 80

## AKK110P1

<211> 424  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 80  
 aacattgttt taattaaaat ttacccctcc tgtagcaatg acatcagaca gacttgcccc 60  
 agtagttcca gatttgacag cccaagagac caacagactt gaacgaacta gttcttttgt 120  
 cgatttggca attcgggatg gagttccata tccacctagg cctgcaatta ataattgttc 180  
 tccataacctg aatatgttga ctggaacgtt ttctgtacca aatgtaaatc agtacacggg 240  
 tgcaataggt ccttatcgac cagcaaatcc tgtttatact tattatagct ataaatgcta 300  
 ttttccgtat agaaattatc gaggctacac actgacggat gcttactggg acgaccgtta 360  
 ttattatatt tcgccaatat acaaacggtc aatgttccca attagattcc ggcattctga 420  
 ctac 424

<210> 81  
 <211> 89  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 81  
 attatccaca cacctattgg agctaccctt accaaggaaa atggtacgac tatgacaatc 60  
 caacanatta ccgcccattc ttgaccca 89

<210> 82  
 <211> 168  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 82  
 tttttttttt taaaatttat tcattaacaa atgaccttaa cagataaaaac ttaacagtca 60  
 aaagacaaca taatttccaa ctttttcaat attatccttt ttaacggttt gattttgcaa 120  
 ctcgctccaa ttcgtccttc ttcttgatag catatgaatt gctcgaa 168

<210> 83  
 <211> 67  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 83  
 aattcatcag ccagacattc agcaattgtt ttgatattac ggaaagaagc ttcacgagac 60  
 ccagtac 67

<210> 84  
 <211> 42  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 84  
 taacacgacg aagaggcgaa acatcaacag cctgacgacg aa 42

<210> 85  
 <211> 429  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 85  
 tatacagagta gaatcctccc gtggtcctcc attaataaca gcgccaacaa gtatttgaac 60  
 tggattctct ccagtcaaaa tatgtataat ttcaaaagcg tgcttcacaa tccgaacagc 120  
 catcaacttt ttaccattgt tacgtccatg catcatcatc gaacaaacca aacgttcaac 180  
 aatcggacaa tgagcctttc gaaaacgttt gatttgatat cgaccagcac tgtgcggcaa 240  
 atatttggcc gatitgtctt taacagcaat ataatccact aaagaagcat cattaacttc 300  
 gatatcgctt aaagaccatt taccaaacaa ttttaatttc ggaaaatcaa ttgtagtcac 360  
 ttgcatatcc ccttgtccac caggaaacac agttgcgccc caattatcat cagcgggtaa 420

accatctcc AKK110P1 429

<210> 86  
 <211> 435  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 86  
 tttgagtttt taaaaagtac atactatttta atttttaaca aattattttg atcaatttta 60  
 aattttcttt tcatcatttt ttaatttaaa aaacatttta acaaattaca agaacaacaa 120  
 acataattgt ctctttttta ttataaaatt taaagtttaa taagttttta aacattctcg 180  
 actggagtac gtgtacttag tgttttagaa aaggcaaaat tagtttggtg gtttgaagag 240  
 acaaattctt ttgcacaagt agcgagaaga tatcgagcag aatttggaat ggaaccccca 300  
 catatggatt tagttaaaaa attacatcaa cgttttctca atactgggtc tgtttctaat 360  
 ggaaatactg aacattttga agttaatcca acaatggaaa catcgacatc ctcaacagag 420  
 ggtgtagcag atccg 435

<210> 87  
 <211> 501  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 87  
 gttttttttt ttttttttta aacaaaatat cgagtcttta taagacaaaa ataaaagaca 60  
 aaagcaattt agttttatca ataaaattaa aaatagtc aaatagtc tgtctcggtt cactcattag 120  
 atttggtggc cttaaagagg ccgtttgggt ttggttggtg tacttcagct gcctccacc 180  
 aattgttctt tagccaccaa atccgtaaag agtacgtcct tggcgtttca acgcatagac 240  
 gacgtccatg gctgtgaccg tctttctctt ggcgtgtacg caataagtta ccgcgtcgcg 300  
 gatcacattt tcaagggaaga ctttcagaac acctcgagtc tcctcgtaaa tgagcccgga 360  
 aatacgtttc actccaccac gacgtgccaa tcgccggatt gccggtttgg tgataccttg 420  
 gatgatatca cgcaagactt ttcggtggcg cttagcgctt ccctttccaa gtccctttcc 480  
 gccttttact cgtccggaca t 501

<210> 88  
 <211> 270  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 88  
 ggaagtgtgt ttaagataaa tggatgatta gaaataaaaa tgaattgatt aaaaattacg 60  
 ttagaataat aatggaatat ataaaaataa attggatgat ttaataaaaa aaaaaagag 120  
 agaactagtc tcgagttttt tttttttttt tttttaanaa ttaacaattt atctcatttt 180  
 cctcttccat gaaaattaac aaaaagacga caacttaatc ccataattaa catcattttt 240  
 aagcttcagt cggcatgctt cgaataatgt 270

<210> 89  
 <211> 286  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 89  
 caagcggttc ccaactcaat gttgttgcca tgatactcgt gaacaccagt tctcgccaac 60  
 atagaatagt actcaatctc actgcgtcta aggcctggag tattattcga aataataaca 120  
 agtttagcct ttccagaacg aagagtcctc aacgtctgct ttagacccaa acaatacttg 180  
 ccgattttgg taacctggc gagacgagca ttgatatttt cigtggactt tttctgtttt 240  
 ccaacaacca ttgtaacgca aaattaaaat ctctttttta acaaat 286

<210> 90  
 <211> 391  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 90

## AKK110P1

```

agatatgaca tcagacagac ttggccagat agttccagat ttgaccagcc aagagacca 60
tagacttgaa cgaactagtt ctttggttga ttttagcaatt cgggatggag ttccatatcc 120
tcctaggcct gcaattaaca atgttcctcc atacctgaat atgttgactc gaacattttc 180
tgtaccaa atgaaatcagt acacgggtgc aataggctct tatcgaccag taaatcctgt 240
ctatacttat tatagctata aatgctatct tccgtataga aactatcgag gctacacatt 300
gacggatgct tatttggtacg accgttatta ttatttttcg cctatatata aacgggtcaat 360
gtttccaatt agattccggc actctgacta c 391

```

<210> 91  
 <211> 131  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 91
attatccaca cacctattgg agctaccctt accaaggaaa atggtatgac tatgataatc 60
caacaaatta cgcgccgttc ttcgaccac gcatcagcgc atcatattca agaccttatg 120
attacacatc a 131

```

<210> 92  
 <211> 571  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 92
ttggtgagac aacaaaaaaa ttttatttat tttttaacaa cagaaaaata tacttttta 60
tttttaatat ttttccatga ttcaacagcc atactttcct ctttttaata cttcttaaac 120
cctcaaaaaa ttcatttatt gacgaccagc agcagggtgt tgctgctgtt gttgaccacc 180
acccccttgc gcttgacctt gctgttgctg tcccttcacg tcaacaggca aatigagttg 240
caaataatca accatctcct tagtctcttg atcaacacta atagttggat gttgagaagc 300
atcaagatag gaaacttctg gaacccaatt atcacgacgc tcacgctctt cttcttgcaa 360
tttaatataga attccacgaa caggctcctt ttcgatacgc ttcatacaat gggtaataaa 420
accagcaatt tgattacgca tccgtttgct aggaataaca gcaatttcct cacaatttcg 480
tttgttcaca tgaaaatcat aagtcaagcg tgtataatat ttgtcaataa taacacgaga 540
tgctttcttg acagttttga gagaaccgat t 571

```

<210> 93  
 <211> 671  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 93
tttgagaatt taacttttct aacaaaaact tttatttttg tctttgatgt ctactcaagt 60
accaatacgc gtgctggtta ctggagcagc tggtcagatt gggtattcct tgggtattca 120
aattgcaaag ggtgatgttt ttggaaagga aacgcccatt gttctggtta tggttgatat 180
tcctccaatg gccgaagtgc ttaaaggagt ggaacttgaa ctttacgatt gtgccttggc 240
gaatcttata gctgtcgagc cagtcacgac tgaagaggca gcgttcaaag acattgatta 300
tgcttttctt gttggtgcaa tgcctcgaaa ggaaggaaag gaacgaaagg atttacttgc 360
tgctaattgtg aaaatattta aatcgcaagg attggctcta gcaaaatatt caaagccaac 420
tgtaaggtt ctggttggtg gaaatccagc aaatacaaat gcttttattt gtgcaaaaata 480
cgacagagat aaaattccag caaagaatgt cagcgctatg actcgtcttg accataaccg 540
tgcaattgcc caaatagctg ctcgttggtg ggttgactgt ggatctgtga agaaagtat 600
aatttgggga aatcattcaa gtacccaatt tcctgatgtt aaacatgcta aagtaattaa 660
aggtggcacg g 671

```

<210> 94  
 <211> 289  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 94
ggctgtaaat gatgtgccgt ggatacagaa tgaatttatt tcgaccgtcc aaaagcgcgg 60
agctgttatt atcgaaaaac gcaaaactgt cagcgcaatg tcggcagcaa agggcgcatg 120
tgatcacatt catgattggc acttggaac aaaagatggc gattgggttt ctatggcctg 180
tccttccgat ggttcttatg gaattccgga aggtttgatc ttctcatttc caattacaat 240

```

## AKK110P1

tgatgcanaa acgctgtgact ggaaaattgt acaaagatta gaactcgat

289

<210> 95  
 <211> 262  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 95  
 aatttaactt ttctaacc aaacttttatt tttgtctttg atgtctactc aagtaccgat 60  
 acgctgtctg gttactggag cagctgggtca gattgggtat tctttgggta ttcaaattgc 120  
 aaagggagat gttttcggga aagaacgcc catcgttctg gtaatgttg atattcctcc 180  
 aatggccgaa gtgcttaaaag gagtggaaact tgaactttac gattgtgcct tggcaaatct 240  
 tatagctgtc gagccagtca cg 262

<210> 96  
 <211> 323  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 96  
 aagacattga ctatgctttt cttgttggtg caatgcctcg aaaagaagga atggaacgaa 60  
 aggtattact tgctgctaatt gtaaaaatat ttaaactcgca aggaactggct ctacgcaaat 120  
 attcaaagcc aactgttaag gttctgggtt tgggaaatcc agcagatata aatgctttta 180  
 tttgtgcaaa atatgcagca gaaaaaattc cgacaaagaa tttcagcgct atgactcgct 240  
 ttgaccataa ccgtgcaatt gcccaaatag ctgctcggtg tgtggttgac tgtgggtctg 300  
 tcaagatagt tataatgtgg gga 323

<210> 97  
 <211> 717  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 97  
 aatattttta acaaacgatg taacagaaaa acaaagtttt ttaacaaat tttcttgaac 60  
 cttatttttt ttcaaaacat ttttttattt aaattttaaact ctctcttcat ttctcttaa 120  
 cactttcttg aactggaggt tcataagcat ctggacgact ttcaataact tctcacttg 180  
 ctgtagttat agcaacttgt ccaccaccac ttccagcacc ctctccatgc atatccaaa 240  
 gttttccaag ttcaattttt gggttttttca aaattttttac ttttcgaata taaacgtctt 300  
 gaagtggata gaaataagaa caagactttt caatgtcttt tccaatagaa tcaggaatta 360  
 atttgctgac aacttcttta agatcgcatg aagaaacctc gcgatgaata atctcaacca 420  
 tcctagcacg aatttgacgc acttgagacg attttgcata actagtcttt ttcaacttgg 480  
 ttggagcttt ctttgtgaag ccaatacaga acaatcgaa gcaataacca tcagtgtgtt 540  
 tgacagcaac atttgcttca attaaagtat gccacttttt gacaatagaa caaagcttgt 600  
 ctcgagttaa agtcattcca tggaaattgg tcaacaaaac tttgccttga acctcttcac 660  
 aaataagtcg aaatttgcg aagtcagctt cgggtgtgtt cagatcacca agagaaa 717

<210> 98  
 <211> 758  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 98  
 gacaagttaa accttgtgtg actttatcta tattcttgtc taaataattc taacaaattg 60  
 taacaacaaa caaaaatggg cgagcaagac aaaaagaag ctggcgccgg cgatgggtggc 120  
 aaaaagaagg atggcttcga tgccaaaaag tttgcgattg atttggtctc tggaggaact 180  
 gccgtgcggg tttctaagac ggctgtggcg cctattgaac gtgtcaagtt gtgtctacag 240  
 gttcaagacg cttctcagca catcgtgcc gataaacgct ataaaggaat aattgatgtg 300  
 cttgttcgtg tgcccaaaga acagggagtc cttgcttttt ggctgtggtaa tttggctaac 360  
 gtgatccgtt actttccaac gcaagctctc aactttgcgt tcaaggacac ttacaagagg 420  
 atcttcatgg aagggtgtga caagaacaaa cagtttggca aattcttttt gatgatgtg 480  
 tttatgagca aaaatttcct tgggtggaaat agacctaaca gttgaagagt atcttgccct 540  
 ctgtgatacg tatacaacac tctcttcaat tggagattca atgttgagtg gagatgctga 600  
 tagtaatcct ctgttacaat cacttaacaa ctcaatcaat tccaatgcca ctgctcagaa 660  
 ttataactcc tcaacaattg gccgaagcta aaaactacgt ttcaacatgc tacagctact 720



AKK110P1

tcaaaatcga aacagattgt tttaaactgt tgaaattt

758

<210> 99  
 <211> 154  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 99  
 ttgagttcgt tggcacattt gttgtgttac aaaacgaaaa ttattgggaa cgggtttcag 60  
 tgcctattct cgcagggtat tggcacttca cacatttgta ccaataacaa cgttaccgt 120  
 tataatcaaa ctgttcctca aagttatgcc catt 154

<210> 100  
 <211> 125  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 100  
 ttcagaatac tcaaggtctt atattcgttg ttgatagtaa cgacaaagag cgtattgttg 60  
 aagctcgtga ggaattgatg cgtatgttgt ctgaagacga acttcgcgat tctgtactcc 120  
 tcgta 125

<210> 101  
 <211> 219  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 101  
 cttgccgaat gctatgaacg ctgctgaact tacagacaaa cttggacttc acacgctgag 60  
 aaatcgtaac tggatatatc aggctacttg tgccacttca ggagatgggt tgtatgaagg 120  
 tttggactgg ttgagtaacc aattgaagaa tcaagggttaa atgagtctaa ataaaaatgg 180  
 agaggggaaa gaggagaggt taatttttta aggaaaaaa 219

<210> 102  
 <211> 473  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 102  
 gttttttttt ttttttttta aattccaagt tttcttccaa atgagagaat agggagaatg 60  
 atgggggaaa aaataggagc aagccaaaaa gccaaaaaaa aatttttttt ttaaattgatt 120  
 ttgtaaaatg tgtgaaaagg tgtgtgtcaa ttgtagagtc aaatgtcgtt gccttccttc 180  
 cactaaaatt tctctttcct ttcttttctc ttctaaaaatt cttcaaaagt cgatccaacg 240  
 aaatttcagc ctccctctgga tattccaact cccaaatacg cttcaaatgt ttgcctttaa 300  
 cgtcacgagg agtaccaaact ccagtcatca acttttgaga gtctccctta ttccaaccgg 360  
 cctgggatgg aattatcgtt tctgacttct tcatatcttc atatggaagt tcgccagact 420  
 ccgctcgtga tgttggtgtc cttggcgttc caaaacctgt catgcccgtc tgc 473

<210> 103  
 <211> 114  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 103  
 ttggaccgtt aggattgtcg ccaaagaaaa ttggagaaga cattgcaaag gcaacacaag 60  
 actggaaagg cttaaagggt acttgcaaat tgactatcca aaaccgaatt gcc 114

<210> 104  
 <211> 255  
 <212> DNA  
 <213> Meloidogyne incognita

## AKK110P1

&lt;400&gt; 104

```

ccgcttctcg aattgtgaag gaattgaagg aacctcaccg agaccgcaaa aaagtcaaac 60
acgtaaaaca cagtggaaat ttgacgatcg agcaaattat caacattgca cggcaaatgc 120
gacctcgttc aatggcgaaa aaaattggaa gggactgtta aggaaattct tggcactgca 180
caatctgttg ggtgtactgt tgatggacaa catccacatg atattgttga tgcaatccga 240
agtgggaaaa ttgaa                                     255

```

&lt;210&gt; 105

&lt;211&gt; 571

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 105

```

tttttttttt tttttttttt tgtcaacaat aaattttactc agaaaaatca ttttaacaatt 60
taacacacat ttttaattcc ttaatactcc aaaaaacttc tcttctttat tccctcttat 120
tctccaatt catttaaagt ttcagttttg tgcggcgcca atgacgacgt tttgcattat 180
agcgtatagc actgccagtt ttcattcgaa cccattgctg cagcggcgca ttttgtttag 240
cagccttagc cagcttgctc ttgataataa acgttttgtg tgcagccatt aaattgttga 300
ctttatccaa aattgttttt ttgaaggcaa taaacaaatt taatttttct gctcaacaag 360
tccatagcag ctcatctggt caacaatctc cctcatgctc ctcatgtctc agcgtttcct 420
cttatgaatg tcaaaaaacag cagcaacaac ccccgagcaga accttgtgga ctttcttttg 480
aagtctcatc atctggtcat tcaacaacaa cctttccatc tccatgttct ttattacccc 540
ctccctcttc ttacatctct ataatcatc g                                     571

```

&lt;210&gt; 106

&lt;211&gt; 235

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 106

```

tgctttattt tcaattcttc aaccaaaaaat taaatcttcc cttattttta ttacaattcc 60
aatttttagc gcattagccc caactacttt agctgctaata aaaattgttt atgaggatgg 120
agatagtgat ggacttgata tggctaaaag tatttttaaat tgaataaagg aaaaagaagc 180
atttttaaaga aaattagatg gaaatgctga agaaagaaaa aaattattta ttttt 235

```

&lt;210&gt; 107

&lt;211&gt; 702

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 107

```

ttttttcaaa aaataattcg aattttgttc ttttttatTT tgctacaaat aaaattttaaa 60
tttgaaaaaa aaaaaaaaaa aaaaaaaac tcgagaagaa atccttgccg aaattgacgg 120
cttcaaaatt gagagtatc aacgtttctt cgatatgttt gaccgtggaa agaattggcta 180
tattatggct actcaaattg gggttaattat gaatgctatg gaacaagatt ttgatgaaaa 240
aactcttcg gaaattaatcc gaaaattcga cgcagacggc agcggcaaaa tcgaattcga 300
cgaattctgc gctttggtat acactgtggc gaatactgta gacaaggaca ctttgcgga 360
agaattgaga gaagcttttc gtctctttga caaagagggc aatggttaca tctctcgcc 420
aacactcaaa ggattacttc acgaaatcgc cccagacctc agcgataaag acttggatgc 480
cgcagtagac gagatcgacg aagacggaag cggaaaaatt gaatttgaag aattttggga 540
gttaatggct ggagagactg attgaaattt taattagaat gactagaaaa ttgactaaaa 600
tattttgcca tttaaattttg gaaaagtcca aaaattgcct ttctgagaat ttttattttt 660
aacgtctaaa taatgaataa aatggatata aaaaaaaaaa aa                                     702

```

&lt;210&gt; 108

&lt;211&gt; 423

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 108

```

aaaatttaaaa taaaagacaa acaataaat ataaatttaa taaataatat ttaaataaac 60
acacaaataa actctccaaa cataattttt ttaaatttta ataactttt gtcccatTTg 120
agaaagaaaa tgccaaagga gatgaagaac ttgttgaaga aaaaagtTca aaaatatcaa 180
ctcctccatt tgcgtcaca ttttctttca ttattccatt tgttgaagc tcagtaactg 240

```

## AKK110P1

```

cccccaattgt tgtttagtc catggagaga aagcactttc ccatttcgaa aatgttgaac 300
caaattgggtc aaattgttgc tgttgttgac ctggaagttc gttagaaaca gaacgaaata 360
aattatgagg ttgttgttgc tcctgacgtt tttgattgtc tggagctggg tgaggatcac 420
caa                                                423

```

<210> 109  
 <211> 994  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 109
ttttattttt tatttgaaaa taatcatcac attataatta atgggaaaaa gacaaaaaat 60
tagaacagggt gctggcgatc ttgtcacaaac ccctggacct cttcataaac aaattgaaaag 120
gtcaaaaacta gccaagccga aattcaagcc tttaaaacgt tcaagagaag agcaaaaaga 180
tgaaattgaa cttgtcgatc catcgttaaa gggcaaaatt attattaaag caaacaaaaa 240
attggaaaaa gatgttgtgt tcaatgagga tggagaatct gataattctg aagaaattga 300
agaagaagaa gaagacggca atgaaaagtt ggatgttgat caattagtat caaaacattt 360
ggaagattta gatgaactaa aattggatga tggcgttgaa aatgtgcgaa agataataac 420
gaaattcaga taaaaataac aaagaaagtg ttataaataa agctgagttt gccgatatcg 480
acccaaaaat tgttgatctt tttacagaaa ttggtcaagt tttaaagaaa tatagaagtg 540
gacgtattcc caaagctttt aaagtatttc caactttggt tgattgggag aaaattatcg 600
aattaaactcg ccagatgat tggtcggcag ctgcaatgtt acatgctacc aaaaattttg 660
cttcaactgc taccctact caatgccaaa ggttttataa ttgattttg ttgccacgta 720
ttcgagatga tattgacgga ttaaaaaaatt acatttccat atgtatcaat gcttatttaa 780
agcattgttc aaaccagctg catttttcaa aggaatcctt ttgccgcttt gcaaatcgaa 840
caatttttct cttcgagaag ctgttgrtct tgcttctatg cttcgtaaag cctccatccc 900
tcaattacac gcggccgcag cattgttgag tatttcttgc ttagaatata cttcttcaag 960
ggcttatatc cttcaagcat tgatagaaaa gaat                                                994

```

<210> 110  
 <211> 476  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 110
tttaaacact taaaaatacc ttcaaattta ttttagaacc tttttgccat taaaaaaaat 60
tttatttcga aaaaatggct gagaatatag aagaaatcct tgccgaaatt gacggctctc 120
aaattgagga gtatcaacgt ttcttcgata tgtttgaccg tggaaagaat ggctatatta 180
tggccactca aattggggta attatgaatg ctatggaaca agattttgat gaaaaaactc 240
ttcgaaaatt aatccgaaaa ttcgacgcag acggcagcgg caaaatcgaa ttgcgcaaat 300
tctgcgcctt ggtatacact gtggcgaaata ctgtagataa ggacactttg cggaaagaat 360
tgagagaagc ttttcgtctc ttcgacaagg agggtaatgg ttacatctct cgtccaacac 420
tcaaaggatt actccacgaa atcgccccag acctcagcga taaagacttg gatgcc 476

```

<210> 111  
 <211> 189  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 111
cgaagacgga agcggaaaaa ttgaatttga agaattttgg gaattaatgg ctggagagac 60
tgattgaaat ttttaattaga gatgaataaa aaattaacta aaatatattg ccataaaaatt 120
ttggaaagtg ccaaaaattg cttttttgag aatttttatt tttaacgtct aaataatgaa 180
taaattggat                                                189

```

<210> 112  
 <211> 164  
 <212> DNA  
 <213> Meloidogyne incognita

```

<400> 112
ttgaggaaat ttaatttttt aaacaaatat aataattacc aaacaacaaa aaagaatccc 60
aaaaacaaca tttttaaatc aaatgacaga catatatattg caataacgat gtgtggattt 120
tctttttttt taaataatta acatcttaag cctgctattt cttc                                                164

```

## AKK110P1

<210> 113  
 <211> 539  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 113  
 cagctttctg cgcagatttg gtaacctttc caccagcttc gaccttctcg acggccttga 60  
 taacaccaac agccacagtt tgacgcatgt cacgaacggc gaagcgtcca agaggagcgt 120  
 agtcagtaaa agcctcaaca cacattgggt tggttggaat taagtcgaca ataccagcat 180  
 ctccagtcct caaagccttt ggattgtctt caaccttctt tccagttcga cggtcgacct 240  
 tctctttaag ctccagcgaac ttgcaagcaa tgtgagcagt gtgacagtca agaacaggcg 300  
 tgtagccagc agcaatctgc ccaggatggg tcatgatgat aacctgagca gtgaattgct 360  
 tggctctcctt tgctgggtca ttcatagagt cagaagtgcac tgaaccacgt cggatgtcct 420  
 tgacagagat gttcttaacg ttaaattcaa cattgtcttc aggaacagct tcagggagag 480  
 actcgtggtg catctcaaca gatttaactt cagtagaaat tccttcagga gcaaaggta 539

<210> 114  
 <211> 314  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 114  
 gtttttaatt ttagaaaatg tctacagaaa cagaaaagga tttagaacgt tgggaggatg 60  
 tccgtcgatt tactgagatt gggtcttcta aatttgccca tcccgctttt gttccaagcc 120  
 cggagaatct tgaaagagta aggaaatgtc cagttttggg tgttggtgct ggtgggcttg 180  
 gatgtgaaat tttgaaaaat ttggccttat caggatttca aaatattgaa gttattgata 240  
 tggacacaat tgacctttca aatctcaaca gacagttttt gtttcgtgaa cacgatgttg 300  
 gcttatacaa agca 314

<210> 115  
 <211> 200  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 115  
 ttcgaagacg tgttaaagga tgtcgtctta ctgcacataa ttgtaaaata caagataaag 60  
 gacttgactt ttatgggcaa ttttcaatta taatttggg actagattct attgatgctc 120  
 gaagatgggt aaacgccaca gtgtgttctt tggtcgaatt tgacgaagaa aacaagccac 180  
 ggccaggcac aattattcca 200

<210> 116  
 <211> 471  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 116  
 tttggtcgaa aaaagactgc tactgctgtg gcatattcca aaaagggaaa aggattaatc 60  
 aagggcaatg gccgtccttt agaatttttg caacctgaaa ttcttcgtat taagctacaa 120  
 gagccattgt tgattgtagg aaaggacaaa tttgctggaa tggatattcg catccgtgtc 180  
 aaagggtggtg gtcattgtgc acaaatttat gcaattcgac agtcaattgc taaagttttg 240  
 gtggcctatt accagaaaaa cgtggatgag caaagcaaga aagaattgaa ggatcaactt 300  
 gttgcttatg atcgtaattt gcttgttgcc gatccgagac gtcacgagcc aaagaagttt 360  
 ggaggacctg gtgctcgtgc tcgttatcag aaatcttatc gttaagaagt atgaaattat 420  
 aaaattgtgt gttacgaatt aattgttatt ttgttgggat aaatntgaat a 471

<210> 117  
 <211> 593  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 117  
 gaattcaaaa aatattaaaa ttgtttaata taatttctaa aatgaagcca aaggttggaa 60

## AKK110P1

```

ttaacggatt tggacgtatt ggacgtcttg ccctgcgtgc agcggtcgag aaggatactg 120
tccaagttgt ggctgtcaat gaccggttca ttgatcttga ctatatggtc tatatgttta 180
actatgattc caccacgga cgctttaaag gaaagattca agcaagcaat ggaaatttgg 240
tagttgagaa ggagggaag tctactcata ctatcaaagt tttcaacttc aaagaacctg 300
aaaagattga ctgggcaggt tctggtgctg attttgttat tgagtcgact ggagttttta 360
ctactaccga gaaagcttct gctcacttga agggcggagc caagaaagtg gttatctccg 420
ctccatctgc tgatgctcca atgtttgtgg ttggtgttaa tgaggacaaa tatgatcctt 480
ccaagcatca tatcattagt aatgcttcct gcaactactaa ttgtcttgct cctcttgcca 540
aggttataaa tgacgagttt ggcataattg aaagttgaat gactactgga cac 593

```

&lt;210&gt; 118

&lt;211&gt; 576

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 118

```

gaattccgag tttttttttt ttttttttaa aacaaaaatt aaaagattta tcgccatcct 60
ttgccagcca ttgtcccgcc atttttttgt gcacaataaa tttttttgta atttttgggg 120
tgagggggaa gtaaaatgaa agaagggaga gagatatgaa ttggaggitt ttttgttaaa 180
ataaaatttt ttttcttgaa aattcttccc gtttctgagc ttttctgctt ttttcaatt 240
ttcgtttgtc gaaatactaa actttacaat ttggttaggt tctatttgtg aaacataaat 300
atctccatta tcgctgattg caagggcatg ggcgttttcg agaccctttg caaagctatt 360
agcccttcct gtgttcatat ccattacgaa aacttgggat tctaattgac tgccttgatc 420
ttgattggtg acgccgacga ggaagtgttc tttctctcgg atagcaaaga ctgcaccaat 480
attttcagcc tttgtgaaga aagtgcctgt ggggacgtaa gcacgtctat gttggtgttg 540
agcgccttct aatccagcag aaaagcattg aatacag

```

&lt;210&gt; 119

&lt;211&gt; 559

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 119

```

acgcagagta agttgagatc ttcaataagg gttagagagt gtggtacgag gaattctcca 60
tttttgggtg ttctactgga gtcaggcttc ccaaattgac tgagcaattt cccatccttg 120
tcaaacttca ttattcggct attacagtaa ccatctgcca cgaaaaactc tcctgtactg 180
gcaatagcaa cgtctgtagg tttgcaaaaa ttgttgcat ctgtccctgg aacaagcttt 240
tcgccccaac tcataattaa tttaaaatcc ttgtcaagtt tgtggacttg atgacttcca 300
acgtcagtaa cccaactatt gccgtgggca tcgattgtta gtccatgagg catgtaaaac 360
atgctttttc cgtattcttc caagactgcc cctgattccg tgtctataac agcaattgtt 420
gtgtttgaaa tgatgccag ggatctgttt aggtggttgt tctcatcaaa cgaaaattca 480
tcccaaactc tgtcagatcg gtgaaaaaga acaagtcgat tcaatggatc caatgcaata 540
cccggagctt gcccaatat

```

&lt;210&gt; 120

&lt;211&gt; 366

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 120

```

tttaagaatt ttttaaaaat taaaacttgg actagatttt aataaaatgt cagctccacg 60
tagtgttgct agcgggtgtg gtgctgctgt tatgaataag caagcaagta aatacaatga 120
agttgaagga gaactccttc ttaattggat taagaaagtg acaggcgaaa atattgctat 180
aaacggaact agggaaaatt ttgtgaaaca attgaaagat ggaactctgc tctgcaaatt 240
tgctaacaaa attgtgccaa attcaatcac aaaggcacag gcaaaaccga acagcacatt 300
ccaatatatg agcaatttgg agctgttctt aacatttatt tcaagccaag gagtccctag 366
ggagga

```

&lt;210&gt; 121

&lt;211&gt; 661

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 121

AKK110P1

ttagttgaat	ctcgtgacct	ctactctggt	tgtatgacat	taaattctct	tggccgcatt	60
ttggaacgtc	aaggaaaaac	tcatccagag	cagggttaagt	cgtcagaaat	tcttaatttg	120
ggtactggag	accaagtgcg	ccttcgtggt	taaagatggg	aaattgaaag	aatttttggt	180
aaacataata	aaaagacatt	ttatggcaat	aaaaaatgt	caaaaaagct	tgtcttttaa	240
atattttggc	aaaacatttt	actttcacaa	aattttaaaa	taaatttatg	aagattgttc	300
cgtcactttc	atcattttccg	atcgaccttt	gttggtttct	aagttcgttg	gccaaagaaa	360
ggatatgtaa	aattgaatta	tgaataaaaa	taaatcactc	aatcagaggc	attgttagtc	420
tctcacttcc	tcctcttttac	ccattggcta	accagcttta	aggatttttt	ccataagttc	480
aaggtgtacg	taaatacgaat	accgactgtg	gtatcttaat	ttttccatga	aattctccaa	540
taaaaaaaaa	ttttttttat	tttttttcca	taatgctatc	tatatatttt	gcttttaatc	600
ttttttggct	atcaggcctt	aaaatagtaa	atatacttat	attaatattt	tatttccttt	660
a						661

<210> 122  
 <211> 173  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 122  
 ggagagtttt tctgtggcaga tggttactgt aatagtcgaa taatgaagtt tgacaaggat 60  
 gggaaattgc tcagtcaatt tgggaagcct gactccagtg aaacacccaa aaatggagaa 120  
 ttccttgtag cacactctct aaccctcatt gaagatctca acttactttg tgt 173

<210> 123  
 <211> 584  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 123  
 cgcattcaat gcttttctgc tggattagaa ggcgctcaac accaacatag acgtgcttac 60  
 gtccccacag gcactttctt cacaagggtt gaaaatattg ggcgagtcct tgctatccga 120  
 gagaaagaac acttcctcgt cggcgctcacc aatcaagatc agggcagtca attagaatcc 180  
 caagttttcg taatggatat gaacacagga agggctaata gctttgctaa gggcttagaa 240  
 aacgcccatt cccttgcaat cagcgataat ggagatattt atgtttcaca aatagaaccc 300  
 aaccaaatgg taaaattttag tatttcgaca aacgaaaatt gagaaaaaaa aaaaaaaagc 360  
 tcagaaacgg gaagaatttt caagaaaaaa tttttttacc aaacaaaaaa cctccaattc 420  
 atatctctcc cttctttcat tttctctccc cttctctccc aaaaattaca aaaaatttta 480  
 ttgtgcacaa aaaaatgggc gggcgggcga atggctgggc aaaggatggc gataaatctt 540  
 ttaatttttg aaaaaaaaaa aaagaattcg aattatatgg ccta 584

<210> 124  
 <211> 650  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 124  
 gtttaagaca attaaaacgt ttattttcta caatcaaaac aaatatggct gttcctcccg 60  
 atgttatcga gaagatcgag gctgggtaca aaaagttgca ggaggcaccg gagtgaagt 120  
 ctcttctcaa gaagtacttc acgaaggaa tttatggacca gtgtaaaggg ctcaaaacta 180  
 agcttggtgc gaacttgctt gatgtgatcc actctggagt tgcgaatctc gatagcgggtg 240  
 ttgggtgtta tgcgcctgat gctgagtctt acactctctt caaacccgctt ttgacccga 300  
 ttattcagga ttaccacaat ggattttggac ctgaccagaa gcagccgcaa actgacttgg 360  
 gtgagggaaa gactcagctt ttgcctgatc tggatcctga gggtaaattc atcaactcga 420  
 ctcgtgttcg atgtgggctt tctcttcagg gatattccgtt caatccgtgc ttgactaaag 480  
 agaattatac ggaaatgcat gacaaagtta aaggggtttt tgagcagctt aagtctgatg 540  
 ctgagcttgg tggcacctat tatcctttgg agggaatgac caaagagggtt caaactcaat 600  
 tgatcaagga tcacttcctc ttcaaagaag gagaccgctt tttgcaagct 650

<210> 125  
 <211> 1013  
 <212> DNA  
 <213> *Meloidogyne incognita*

<400> 125

## AKK110P1

```

tttttttttt tttgatgttt ctaattttttg tgggcaatat ttaatattat ttttaattat 60
taaattttct tctttatttt ttaaaaaatt atttcttaaa tttattcttc tcctcttcgt 120
gttttgatc aaataattaa attttaaatt atttaaacag ctacacgagg cctcagcctc 180
ccccgttgca ttcaaattgg tcggcacggg tggcgatgat aattttattt tttaggtaat 240
tttggtagaga aaatatTTTT aaaggtaata atgtcctttt ggacaattaa aaaaaaactc 300
gaggagagag tgaatatTTT tacaaattat ttgaagagca gccagcctat tgttatcaac 360
aaaaaacctt caaaatgcc aaaaatgatt atgatgagga ggaggcgcca aacgccacga 420
tggaacaaa ggtagcttca ggtggacagc caaaacgctg ttggaaaatg gacattatcc 480
cagctgcgcc agactgatgg tataattcca tcccaggccg gttggaacaa gggagactcc 540
caaaagtga tgaccaattt tggtagctca cgtaacacaa caacaaaat tcgtgctgaa 600
tgccttgctg atgtgcctga agaaattgct cttaaagtc acggtgaagt acgcctcaa 660
tccggtacta accgttttgc ttcgcagaag ggaatgggtg gatttggtag tggacgtgac 720
ttatgcagag aaggagtgtt tgtgagtcaa gaccagccg atttatagcc cctccagaa 780
gagataatcc gtgctagcga ttggaattgtt cgtctccaat ccggtaccaa caaattcgac 840
tcccaaaagg gaatggtcag ctccggtaca aaccgacgcg aaactacaag aatgaaagac 900
accaaacatc cggaatacaa ccacgaagtt aacattgacc aaagcgaat tcctttgcaa 960
tctggtacaa acaaattcgc atcccaaaag ggaatgacca gcttcggtac aaa 1013

```

&lt;210&gt; 126

&lt;211&gt; 80

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 126

```

tggttgacac tgctcaccca gaatacagtc acgaaagcag catcgatcaa acgagcattc 60
cttaccaaat gggatcaaat 80

```

&lt;210&gt; 127

&lt;211&gt; 585

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 127

```

agggaaatgac ttgcttttga cagccacggt gggaggtgct tgacccgagc attagctacc 60
agaaccgtaa atcacaagga atggtccgct tccaatccgg aacaaaccgg gtcgcctcgc 120
aagcgggcat gacaggtttt ggaactccaa ggaacacaa acacgagcgc gactctggcg 180
aacttccata cgaagatatg aagaagtcag aaacgataat tccatcccag gccggttga 240
ataagggaga ctctcaaaag ttgatgactg gatttggtag tcctcgtgac gttaaaggca 300
aacatttgaa gcgatatttg gagttggaat acccagagga ggctgaaatt tcgttggatc 360
gactttaaag gaattttaga agagaagaaa gaaaagagaa atttagtgga aggaaggcaa 420
cgacatttga ctctacaatt gacacacacc ttttcacaca tttaaaaaat acattaaaaa 480
aaaatttttt ttggcttttt ggcttgctcc tattttttcc ccccatcatt ctccctattc 540
tctcatttgg atgcaaaactg gaatttttaa aaaaaaaaaa aaaaa 585

```

&lt;210&gt; 128

&lt;211&gt; 287

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 128

```

catctggaga aacgttgagg caatacatcg ttattggccg taaacttcct acagagaatg 60
agccaaatcc aaaactttac aaaatgcaaa tttttgccag taatcatgtt gttgctaaat 120
cgcgtttctg gtactttact agtatgttgc gtcgtgttaa gaagactaac ggagagattg 180
tttcgtgtca ggaggttttt gaaaagaaga taggctctgt aaagaattat ggaatttggc 240
ttcgttatga ctctcgaacc ggtcatcaca acatgtaccg tgaatac 287

```

&lt;210&gt; 129

&lt;211&gt; 175

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 129

```

gctgtcactc aggcttatcg cgacatgggt gctcgtcatc gtgctcaagc cgatcgaatc 60
caaataatca aggttcaacc gatcaaggct gccgattgca aacgtactgg agttaaacag 120

```

AKK110P1

ttccacaact cttcaatcaa gtttcctttg ccgcatcgtg tgaatgacaa acgtc 175

&lt;210&gt; 130

&lt;211&gt; 599

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 130

acttttgttt	ataatcacat	ttgcattact	ttccgtccat	ccttctttga	gacagaattt	60
aaaggttcac	cttctaagta	aggattgtag	cggctgtatg	attgatgttg	cttttgttgg	120
ggagcaatag	aacgcttgcg	tcgccgaggc	tcctcagccc	tagtaacgtg	aaatttcttt	180
gcaatcatcg	atttgtgtag	tccatttttg	gctaagacct	gttctaagtc	ttgttcatat	240
tgttcagaat	tgctttttga	ttgacagtta	aacatgtgtt	cttggtcaca	aaggcattgc	300
tgattggcct	ggtagctacg	cgagaaatcg	gcggtgttat	caaactcttc	caaacatcca	360
tctcgactgg	agtatcccac	agggcagggg	tttggagggg	cacaatatgc	tggcaaaaca	420
ttgtcactct	taatctcttg	gcggtgtgaa	aattcagatt	ctggatggag	ttgttgggtct	480
ccttcaccgg	cacctcctgt	cataaattta	tgcccaaacg	caatgggccc	ggaagcactt	540
tcaatgtcac	gagaaatcaa	gtcgattaat	tgtgaatgcg	gaaatatagg	ctccccaga	599

&lt;210&gt; 131

&lt;211&gt; 466

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 131

gaagattgga	tttattggcg	ctggaaagat	ggcacaggca	ttggccagag	gactaataaa	60
ttctggacgt	tatccttcac	aaaatttgat	ggctagtgtg	cctaagactg	atgtctcttt	120
attggaggat	tgcaagaggc	ttgggagtaa	tacagcacat	gataatgcac	aagtgtctcg	180
tgaaaatgat	gtggtgatta	tagcaggtaa	accaactatt	gtgtctaaag	ttgcttcgga	240
aattgcacca	gccatccgcc	gagatcatgt	acttatttct	atagcattgg	gcaccacat	300
acgctacatt	gagcagtaat	tgacttcaga	atcccgaatt	gttcgtgtaa	tgccagatac	360
tcctgtagg	ggtaggagca	ggctgctgca	gccatatatc	attgggatca	gcattgtcag	420
gataaggtgat	gcccagatag	ttcaagatct	tctgataacg	ctgggg		466

&lt;210&gt; 132

&lt;211&gt; 266

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 132

atgaaattcg	agttctttgc	atcaaggccc	gtgaaatttt	tctttcgcaa	cctattttgc	60
tggaattgga	agcgccgttg	aagattttgt	gcgatattca	cgggtcaatac	aacgaccttt	120
tgcggttttt	tgaatatgga	ggttttccgc	ctgaagcgaa	ttattttatt	ttgggtgatt	180
atgtggatag	aggaaagcag	agcttgga	cgatttggtt	gctgttggtc	tacaagatca	240
aatcccccca	aaattctttt	tgctga				266

&lt;210&gt; 133

&lt;211&gt; 308

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 133

tctatcaacc	gaatatatgg	attttacgat	gaatgcaaac	gcagattttc	tataaaattg	60
tggaaaacat	ttactgattg	cttcaattgt	ctgccaattg	ctgctgtgat	cgatgagaaa	120
atattttgtt	gccatggagg	tttgtcacca	gattttgcaga	atatggagca	aattcgaaga	180
attatgacac	cgacggatgt	gccagataca	ggcttctctt	gcgaccttct	atggtctgat	240
ccagaccaag	atgtccaagg	attgggagaa	aatgatcgtg	gggtctcttt	cacttttggg	300
ccagatgt						308

&lt;210&gt; 134

&lt;211&gt; 335

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita



## AKK110P1

<400> 134  
 taaatttagt ttcttttctt ccatctcttt ttatgttttg aaagagtgtg ccaaaacaaa 60  
 tggccgcccg tgatggaaga agcaggcaaa attatttaca agaacattca attcctcaac 120  
 tttttgaggg ttaaatgact ggacttatat acaatcaacc aatcgatcct attcaatttt 180  
 tggagaatgc aatagctaaa cttcgaaaaa atcctgatct tccattaaag tgggatactt 240  
 ttataagtgt ttgcctcaa caacagcaac aacaacagac gagaatgaat actggagaaa 300  
 atgcagtttc ttataaacia agcactccta tcgaa 335

<210> 135  
 <211> 506  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 135  
 tttttttttt tttaaaaatc aacagattta ttcaagtgcc tcgggcaaat aacaacaaac 60  
 attccacaaac ataattattat tgaacttttc ctttttataa ctatcaaag gccttctttg 120  
 ttcttgagac ttgtatcacc ttcaaaacat taaaacgaac agttttactc aaaggcctgc 180  
 attcaccgat cgtgacaata tcaccaatag agatatcacg gaaacatggc gaacagtga 240  
 cggacatggt tttgtgacgt ttctcgtatc gacgatattt cggaaacaaag tgcaataat 300  
 cagccgaat gacaattgtg cgctgcattt tgttcttgat aacaacacca gtcaaaatac 360  
 ggccacgaat tgaaacattt ccagtgaag gacacttttt gtcaatataa ttgccttcga 420  
 tagcctcgcg tggagtttta aatcctaacc caacattctt ccaataacga tccttatttt 480  
 tcggctnttt gccaatccct tgcgtc 506

<210> 136  
 <211> 230  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 136  
 aattcctcaa actctgccct ggctgtcctt ctcaaaacga caccctcgct ttattatcac 60  
 ctccagtcac ctacgaaaat tctttgcgag atcaagggag taattcgaca ttatggattc 120  
 tttgttggg ttttaattgt ttatttttgc tactaatttt ctttctaatt gccgcctacc 180  
 tccgttgcg catttttggc tccgccccct acaaaaacca gttccgtcgt 230

<210> 137  
 <211> 216  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 137  
 acaatacac aacaacaaa tcattgttta ccccaaatat ccaaaagttc tcctccaact 60  
 tcaactcgctg ttgtaccaaa ctctactagc tgtaattcgt ccttagctgt gccgttaatt 120  
 tctagtgaat cggaagaaag tgatgaacaa caaaagacgg gggaatggac aaatctaaca 180  
 ttattaatta tttatttcta tgattgtaaa ttgcat 216

<210> 138  
 <211> 395  
 <212> DNA  
 <213> Meloidogyne incognita

<400> 138  
 atgcattcct gaagcaattt tgggtatgga cattgtatgc caagcaaagt ctggtatggg 60  
 gaagacagct gtatttgggt tggcaacact ccaacaattg actccagttg acgggacggg 120  
 ctctgttctc gttatgtgtc acactcgcga acttgctttt caaatttcaa aggaatatga 180  
 aagatttagc aaatatatgc ccggaactaa ggtttcgggt ttctttgggt gtatgccgat 240  
 caagaaggac gaggagactt tggctaagaa cactccgcac attgttgggt gcactccagg 300  
 gcgtctgctg gcgttgggac gtacaggaca attgaagctg aaaaacatca aattcttcgt 360  
 tttagacgaa tgtgacaaaa tgattgggga cgctg 395

<210> 139  
 <211> 591

## AKK110P1

&lt;212&gt; DNA

&lt;213&gt; Meloidogyne incognita

&lt;400&gt; 139

```
gaattcggcg ttgtctcggt gtccacgctc aatttcaccg aaatttttgg ggcaggcgctc 60
ctccacacca aactctgggt cattgacaac cggcactttt atctgggttc agcaaaccatg 120
gactggcagt cacttactga agtcaaggaa atgggtctta tgctgttgaa ctgctcctgt 180
ttggcgtggg aactgagcaa aatatttgcg atttactggc ggattggaca gaatcacaat 240
cgcttgcccg ctgtttggcc agtttattta caatcaaaat tcaacgctca acaccaatg 300
gaaattcatt ttggacctga gccctcgcac acgtacattt cgcactcgcc tgagaagtgt 360
aacccaaagg gcagagaaca cgacctttcg gccatatgct catgcatggg aaaagccaac 420
gaatttggtc gaattgcggt aatggattat attcctgcaa caatttacat gccgaatggt 480
aacaacatat attggccatc gatcgatgac gcgataagaa cggcagctta tcgggggtgtg 540
aaagttgacc tttggtgagt ctgtggcccc atttgaatga acgagcgatt t 591
```